



Field Trip Guide Book - B19

Florence - Italy
August 20-28, 2004

Volume n° 2 - from B16 to B33

32nd INTERNATIONAL GEOLOGICAL CONGRESS

THE RECORD OF MESSINIAN EVENTS IN THE NORTHERN APENNINES FOREDEEP BASINS



Leaders: M. Roveri

Associate Leader: A. Landuzzi

Pre-Congress

B19

The scientific content of this guide is under the total responsibility of the Authors

Published by:

**APAT – Italian Agency for the Environmental Protection and Technical Services - Via Vitaliano
Brancati, 48 - 00144 Roma - Italy**



Series Edit

Luca Guerrieri, Irene Rischia and Leonello Serva (APAT, Roma)

English Desk-copy Editors:

Paul Mazza (Università di Firenze), Jessica Ann Thonn (Università di Firenze), Nathalie Marlène Adams (Università di Firenze), Miriam Friedman (Università di Firenze), Kate Eadie (Freelance independent professional)

Field Trip Committee:

Leonello Serva (APAT, Roma), Alessandro Michetti (Università dell'Insubria, Como), Giulio Pavia (Università di Torino), Raffaele Pignone (Servizio Geologico Regione Emilia-Romagna, Bologna) and Riccardo Polino (CNR, Torino)

Acknowledgments:

The 32nd IGC Organizing Committee is grateful to Roberto Pompili and Elisa Brustia (APAT, Roma) for their collaboration in editing.

Graphic project:

Full snc - Firenze

Layout and press:

Lito Terrazzi srl - Firenze

Volume n° 2 - from B16 to B33



**32nd INTERNATIONAL
GEOLOGICAL CONGRESS**

**THE RECORD OF MESSINIAN
EVENTS IN THE NORTHERN
APENNINES FOREDEEP BASINS**

AUTHORS:

*M. Roveri¹, A. Landuzzi², M.A. Bassetti⁴, S. Lugli³,
V. Manzi¹, F. Ricci Lucchi⁵, G.B. Vai⁵*

¹Dipartimento di Scienze della Terra, Università degli Studi di Parma - Italy

²DISTART, Università degli Studi di Bologna - Italy

³Dipartimento di Scienze della Terra, Università di Modena e Reggio Emilia, Modena - Italy

⁴Departamento de Geología y Paleontología, Facultad de Ciencias, Salamanca - Spain

⁵Dipartimento di Scienze della Terra, Università degli Studi di Bologna -Italy

**Florence - Italy
August 20-28, 2004**

Pre-Congress

B19

Front Cover:

*Panoramic view of the Vena del Gesso
(Gessoso-Solfifera Formation) along the west side
of the Santerno valley at Borgo Tossignano*

The record of Messinian events in the Northern Apennines foredeep basins

Roveri M.*, Landuzzi A.°, Bassetti M.A.***, Lugli S.°, Manzi V.*, Ricci Lucchi F.§, Vai G.B.§

*Dipartimento di Scienze della Terra, Università degli Studi di Parma, Parco Area delle Scienze 157A, 43100 - Parma (Italy)

°DISTART, Università degli Studi di Bologna, Via Risorgimento 5, 40100 - Bologna (Italy)

°Dipartimento di Scienze della Terra, Università di Modena e Reggio Emilia, Largo S. Eufemia 19, 41100 - Modena (Italy)

**Departamento de Geología y Paleontología, Facultad de Ciencias, Placa Merced, 37003 - Salamanca (Spain)

§Dipartimento di Scienze della Terra, Università degli Studi di Bologna, Via Zamboni 67, 40100 - Bologna (Italy)

Introduction

Why a field trip on the Messinian salinity crisis in the Romagna Apennines? We believe there are many good reasons indeed. We are going to introduce in these brief notes only some of the most relevant arguments, hoping the participant at the end of the field trip would have completed their own list with many others.

Firstly, although not adequately recognized by the scientific community, the Northern Apennines keep the most complete, and probably the best, memory of the large amplitude paleoenvironmental changes that affected the Mediterranean area during the Messinian age (i.e. the time interval lasting about 2 million years between 7.2 and 5.33 My) and usually referred to as the “Messinian salinity crisis” (MSC).

Moreover, in no places of the Mediterranean area the fascinating history of the Messinian salinity crisis has left so many indelible signs on both the physical and cultural landscape like in the Northern Apennines area extending from Bologna to Rimini.

Gypsum rocks formed during Messinian events constitute an almost continuous and prominent belt (the so called *Vena del Gesso*) along the Apennines foothills. This rocky belt is characterized by peculiar morphology and ecosystems and has been attractive for man use since the antiquity, providing the base for an important social, artistic and economic development. The large-size, twinned crystal variety of gypsum, known as *selenite*, has been used as building stone since Roman times, and may be even before. A short walk downtown in Bologna may quickly account for this; the basement of many Middle Age buildings, included the famous towers, as well as the ancient walls (IV-VIII century AD) are made up of squared selenite blocks (Fig. 1), many of which were recycled from the Roman theatre. Selenite derived from quarries located in the foothills

of Bologna and more to the east in the Santerno and Lamone valleys (western Romagna). Also from Messinian gypsum-bearing successions, but dominated by the microcrystalline, thin-laminated variety of gypsum known as *balatino*, derives the sulphur mineral extensively exploited till the early 60ties in many small quarries and mines of eastern Romagna, i.e. from the Rabbi to the Marecchia valleys.

The great economic value of gypsum rocks stimulated early geological studies, at least since the beginning of the 18th century. As recently pointed out by Marabini and Vai (2003), Luigi Ferdinando Marsili, an outstanding scientist of the *Istituto delle Scienze* of Bologna, carried out around 1717 what can well be considered the first modern stratigraphic study at a regional scale. In an unpublished manuscript, partially reproduced in Marabini and Vai (2003), he describes in great detail the vertical superposition of gypsum and shale beds in the sulphur mines of eastern Romagna, suggesting correlations and recognizing their close relationships with the selenite rocks occurring to the west and pointing out the occurrence of a continuous gypsum belt from Bologna to Ancona. Enclosed to the Marsili's manuscript, is what can be regarded to as the first geologic map in the world (Fig. 2): an



Fig. 1 – Remnants of the IV-VIII century AD selenite walls, preserved and recycled in the Ghisilardi Fava Palace at Bologna .




Fig. 2 – The 1717 geological map by L.F. Marsili (from Marabini and Vai, 2003).

accurate representation of outcropping gypsum bodies connecting and accounting for a series of aligned sulphur mines in the Forlì and Cesena foothills. During the third and fourth day of this field trip we will run over the Marsili’s outcrops again, this standing in witness of the still unchanged interest of this area through the time. The distinct spatial distribution of selenite and sulphur mines, as already recognized by Marsili, mirrors two different stratigraphic and geologic settings developed during the Messinian in the Romagna Apennines. Roughly speaking, selenite formed through primary precipitation of gypsum from dense brines in shallow-water settings, while extensive sulphur mineralization is related to the diagenetic transformation of clastic gypsum deposits, accumulated in somewhat deeper waters, into limestones. In the field trip area these depositional settings are separated by a tectonic feature oblique to the main Apennine structural trend.

Three centuries after the pioneer work of Marsili, the Messinian outcrops of this area still offer the unique opportunity to observe a complete sedimentary succession developed in both shallow- and deep-water settings, thus allowing the reconstruction of their genetic and stratigraphic relationships.

In fact, contrary to the Apennines, Messinian

successions preserved on Mediterranean continental margins are always reduced and incomplete, being more or less deeply cut by a subaerial erosional surface developed in the time span during which, according to the popular “deep desiccated basin” model (Hsü et al., 1972), the Mediterranean basin almost completely dried up. The lacking part of the story is recorded by those huge salt accumulations buried in the deepest Mediterranean basins whose true nature and stratigraphy still wait to be fully unravelled.

The complete and expanded record of the Apennine will allow us to focus on several, still open questions concerning the different evolutionary stages of the Messinian salinity crisis and, in particular, on a usually overlooked topic, i.e. the role of tectonics in controlling Messinian stratigraphic patterns.

Concluding these introductory notes, we hope the field trip participants will find it a stimulating chance for thorough discussion. For them as well as for those who will use this guidebook to make the field trip by their own, we hope they would appreciate the strong character of this land, of its people and products, formed and developed through the time, in a continuous interlacing of geology, culture and history of science.

Messinian events chronology and modalities

In fact, it has long been recognized that the sedimentary record of Messinian events of the Apennine foredeep basin does not support the deep-desiccation model. No evidence for the desiccation has been ever found; which is usually explained by the particular paleoclimatic and structural context, that would have led to its premature isolation from the other Mediterranean basins, and to the persistence of a deep-water, non marine basin. For this reason the Apennine foredeep has always been considered an anomaly within the general picture and, consequently, often overlooked in the Messinian debate.

The comprehension of the different aspects related to the MSC is a long-lived issue in the scientific debate. Many important questions remain unanswered. The paleogeography and paleoclimatology of the Mediterranean area, the physical and chemical structure of the water column through the Messinian and the possibly active role of biota during the crisis are still poorly understood. The great advance in stratigraphic resolution obtained in the last years after the adoption of astronomical calibrations and physical-stratigraphic concepts, has certainly improved our general knowledge and chances. The Messinian stage is now well time-constrained; the Messinian GSSP has been defined in the Oued




Fig. 3 – Chronology of Messinian events according to Krijgsman et al. (1999a-b).

Akrech section (Morocco) and an age of 7.251 Ma has been assigned to the Tortonian-Messinian boundary (Hilgen et al., 2000); the top of the Messinian corresponds to the Zanclean GSSP, defined in the Eraclea Minoa section (Sicily) with an age of 5.33 Ma (Van Couvering et al., 2000). Although different views actually exist, in recent years a growing consensus has been reached about the chronology of Messinian events proposed by Krijgsman et al. (1999a-b; Fig. 3). This has been made possible by the well-developed cyclical arrangement of Messinian deposits. According to such model, a classical three-fold subdivision of the Messinian stage can be envisaged: 1) pre-evaporitic (7.251-5.96 Ma), characterized by the common occurrence in deep-water settings of organic-rich, laminated deposits, which record sea-bottom low-oxygen conditions related to a progressive reduction of water circulation within the Mediterranean; 2) Lower Evaporites (5.96~5.61 Ma), the first episode of shallow-water evaporites precipitation in marginal basins; 3) Upper Evaporites, or post-evaporitic Lago Mare stage (~5.61-5.33 Ma), showing the widespread development of non-marine deposits with Mollusk, Ostracod and Dinoflagellate assemblages of Paratethyan affinity (Lago Mare facies; Ruggieri, 1967; Iaccarino & Bossio, 1999); the end of the MSC is represented by the

sudden, catastrophic return to fully marine conditions at the base of the Pliocene (Iaccarino et al., 1999). Based on astronomic controlled cyclostratigraphic considerations, the main Messinian events are considered synchronous throughout the Mediterranean area, pointing to a strong external forcing, tectonic and/or climatic. The duration of the various Messinian 'phases' can thus be estimated: 1.2 My for the pre-evaporitic, 0.35 My for the evaporitic and 0.35 My for the post-evaporitic phase. This age-model fits quite well with the two-step MSC proposed by Clauzon et al. (1996), which represents an updated version of the Hsü et al. (1972) deep-desiccated basin model. Such model implies a diachronous deposition of evaporites across marginal and basinal settings at the boundary between the evaporitic and post-evaporitic stages. In marginal settings this boundary is represented by a subaerial erosional surface deeply cutting the continental margins and thought to have been developing during the acme of the MSC, when the Mediterranean sea-level dropped more than 1000 meters and shallow-water evaporite precipitation was shifted toward the deepest parts. The astronomic polarity time scale (APTS) for the Neogene still shows indeed a gap lasting 90 kyr at the base of the post-evaporitic stage (Krijgsman et al., 1999a-b), related to the deep desiccation of the Mediterranean. The dramatic sea-

		marginal basins Mediterranean-type	deep basins Apenninic-type	U.B.S.U.	
ZANC.	5.33	Argille azzurre Fm	Argille azzurre Fm	p-ev ₂	MP
		Cusercoli fm	Cusercoli fm		
MESSINIAN	5.60	hiatus intra-Messinian unconformity	tetto fm +++++ ash layer +++++	5.5 My	MP
			Gessoso-Solfifera (deep-water mb.) resedimented evaporites		
			Gessoso-Solfifera (shallow-water mb.) primary evaporites	euxinic shales	
	5.96	Marnoso-arenacea Fm euxinic shales	Marnoso-arenacea Fm euxinic shales		

Fig. 4 – Chronology of Messinian events in the Apennine foredeep. Modified from Roveri and Manzi (in press).

level drop was also responsible for changing the drainage pattern in the peri-Mediterranean area, leading to the partial refill of desiccated basins with fresh to oligohaline waters of Paratethyan origin (Lago Mare stage). Despite the accurate and unifying chronology of Messinian events suggesting a homogenous development, it has been recently proposed that the MSC evolved with distinct modalities across the different geotectonic settings (Sicily, Apennine foredeep, Andalusia, northwestern Mediterranean; Clauzon et al., 2001). According to this view, the different modalities can be summarized and translated for simplicity into two end-member successions: a **Mediterranean-type** and an **Apenninic-type**, the main difference being the non desiccated character of the Apennine foredeep basin during the whole MSC.

The Apennine foredeep record of Messinian events

With respect to the supposed different Mediterranean and Apenninic modes of MSC evolution, we offer to the discussion a different point of view. The end-member successions basically – and very roughly – correspond to the record of distinct morphostructural settings: the Mediterranean-type to marginal, elevated areas, the Apennine on to basinal deeps.

The true “anomaly” of the Apennine foredeep is that, due to its geodynamic setting and post-Messinian evolution, both modalities actually occur and that the stratigraphic and genetic relationships among them can be clearly defined. In other words, all along the Apennine foredeep system a true Mediterranean-type succession can be recognized in shallow, marginal sub-basins, while an Apenninic-type developed in deeper ones. Based on a careful review of Messinian stratigraphy carried out in the last ten years by several research groups through the integration of surface and subsurface data and the use of a physical stratigraphic approach (Gelati et al., 1987; Rossi et al., 2002; Bassetti et al., 1994; Bassetti, 2000; Roveri et al., 1998; Roveri et al., 2001; Roveri et al., 2003), a geologic and stratigraphic model for the Apennine foredeep Messinian deposits has been reconstructed. Such model is summarized in Figures 4 and 5 and assumes the development of pre-evaporitic and evaporitic stages synchronous and with the same modalities of the other Mediterranean basins. Primary evaporites were deposited only in shallow thrust-top basins whose formation date back to the upper Tortonian, when the ensuing propagation of the Apennine compressive front led to the progressive fragmentation of a larger and deeper foredeep basin. The end of evaporite deposition is coincident with a paroxysmal acme of a regional tectonic phase




Fig. 5 – The geologic-stratigraphic model for the Messinian deposits of the Apennine foredeep (Roveri et al., 2003).

determining the uplift and emergence of the Apennine chain and the concomitant migration of the foredeep depocentres toward the foreland. The intra-Messinian unconformity, a perfect equivalent of the erosional surface cutting Lower Evaporites throughout the Mediterranean, is associated to an angular discordance and hence it is strictly related to a regional tectonic uplift leading to the dismantling and resedimentation of Lower Evaporites in deep basins through large-scale mass-wasting and gravity flows in shelf and slope areas. Tracing down basin the intra-Messinian unconformity is a problem solved placing its correlative conformity at the base of the resedimented evaporites complex (Roveri et al., 1998, 2001). This unit accumulated in topographic lows during the subaerial exposure of uplifted basin margins, thus allowing to bridge the last Messinian gap (Roveri & Manzi, in press). Based on its physical characters, a high-resolution stratigraphic framework for the post-evaporitic successions (discontinuities of different hierarchical rank, stacking pattern and cyclic organization of depositional system, key horizons) has been reconstructed. A second, younger regional unconformity splits the post-evaporitic Lago Mare deposits into two units; the lower one (p-ev₁), only occurring in structural depressions, is composed by resedimented evaporites at the base, overlain by a siliclastic coarsening and shallowing-upward succession; the upper unit (p-ev₂), made up of coarser-grained deposits, has a general transgressive trend and seals all the previously uplifted tectonic structures, generalized subsidence and/or a base level rise, heralding the Zanclean flooding.

The relatively small field trip area has the great advantage to sum-up all the elements of the larger-scale stratigraphic model. This area corresponds to a series of

wedge-top basins (Fig. 5) characterized by different absolute paleobathymetry and subsidence rates. The two main foredeep Messinian depocenters are presently buried below the Po Plain and crop out in Southern Marche and Abruzzi (Laga Basin) and their sedimentary succession can be easily correlated to that developed in wedge-top basins through very distinctive physical elements. The distinction between p-ev₁ and p-ev₂ unit is substantiated by the clear upward transition from high-efficiency to low-efficiency turbidite systems. Moreover, an ash layer of regional extent is an excellent key-bed in the p-ev₁ unit.

The first and second days of the field-trip will be spent in a marginal, Mediterranean-type context, the third and fourth day in a Apenninic-type one. This will give the participants the chance to contrast and correlate the two stratigraphies and, consequently, to assess some basic Messinian items.

Regional geologic setting

The Romagna Apennines, extending from the Sillaro valley to the west, to the Marecchia valley to the east (Fig. 6), is part of the Northern Apennines, a ENE-verging arc, characterized by compression along the external front and extension in the inner western part (i.e. the Tyrrhenian area).

The Apennine chain formed since the late Eocene as a post-collisional fold and thrust belt in the more general context of convergence between the African and Eurasian plates. The Romagna Apennine is characterized by an outcropping succession of early Miocene to Pleistocene siliclastic deposits, overlying buried Mesozoic to Cenozoic carbonates. This sedimentary succession represent the infill of a foredeep basin system actively migrating to the northeast since the Oligocene (Ricci Lucchi, 1986)




Fig. 6 – Simplified structural map of the Northern Apennines.

and formed above the Adria plate in what is defined the Umbria-Marche domain, the lowest structural unit of the Apennine orogenic wedge (for an updated review of the Apennine geology, see Vai and Martini, 2001).

To the west of the Sillaro valley, this unit is covered by the Ligurian nappe (Fig. 6), a chaotic complex of Jurassic to Eocene deep-marine sediments and ophiolitic slabs (Castellarin and Pini, 1989), formed as an accretionary wedge during the late Cretaceous Alpine compressional phases and thrust over the Adria plate, since the Oligocene (Kligfield, 1979; Boccaletti et al., 1990).

The uppermost structural unit of the Romagna Apennine consists of the Langhian to Messinian Marnoso-arenacea Fm, a turbiditic complex more than 3000 m thick (Ricci Lucchi, 1975, 1981), detached from its carbonate basement along a basal thrust, and showing a deformational style dominated by fault propagation folds (Capozzi et al., 1991). Seismic data show that the outermost front of the Apennine thrust belt lies in the subsurface of the Po Plain (Castellarin et al., 1986, Castellarin, 2001), where several ramp anticlines are buried by a thick succession of marine Plio-Pleistocene deposits. The

Romagna Apenninic is split into two sectors (western and eastern) by the Forlì line (FL, Fig. 7), a complex fault zone oblique to the Apenninic trend. The two sectors defined by this feature has a different structural arrangement at surface; the foothills of the western sector shows a gentle N-NE dipping monocline of Messinian to Pleistocene deposits resting above the Marnoso-arenacea Fm., while the same succession is deformed by several thrust-related anticlines with Apenninic trend in the eastern one. The Forlì line played a primary role in the geological evolution of the area at least since the late Tortonian (Ricci Lucchi, 1986; Roveri et al., 2002)

In a NW-SE cross-section, flattened at the Miocene/Pliocene boundary (Fig. 7), dramatic facies and thickness changes within the Messinian succession occur across structural high related to the Forlì line. Such changes finally correspond to the two above mentioned different Messinian stratigraphies whose comparison and correlation is the main aim of the field trip.

The lithostratigraphy of the Langhian to Pliocene sedimentary succession of the Romagna Apennines consists of four formations (Vai, 1988; Fig. 8):

i) the *Marnoso-arenacea Fm* (MA, Langhian-

Messinian), made up of deep-water siliciclastic turbidites, derived from the Alps and, subordinately, from the central Apennines, is a huge clastic wedge, up to 3000 meters thick, filling a large foredeep basin elongated in a NW-SE direction, whose depocenter migrated toward the NE following the propagation of the compressional front. Toward the top thick and laterally continuous turbiditic lobes are

replaced by slope mudstones (*ghioli di letto*) containing minor turbiditic sandstone and chaotic bodies; this is in turn overlain by a unit straddling the Tortonian-Messinian boundary and made up of finely interbedded organic and diatomite-rich laminites and mudstones, informally named *euxinic shales* (upper Tortonian-lower Messinian). Like the coeval Tripoli Fm in Sicily and Spain such deposits




Fig. 7 – Simplified geologic map (above) and cross-section (below) of the Romagna Apennines. The cross-section is flattened at the Miocene/Pliocene boundary to better illustrate the great facies and thickness changes characterizing the Messinian succession across main structural elements (modified from Roveri et al., 2003).




Fig. 8 – Stratigraphic scheme of the Apennine foredeep Tortonian to early Pliocene time interval. Note that lithostratigraphy refers to marginal successions, i.e. those characterized by deposition of primary, shallow-water evaporites during the main evaporative event.

show a well developed cyclical pattern and record the paleoceanographic changes associated to the ensuing MSC.

The euxinic shales span a 1.5 My time interval, characterized by well-defined bio- and magnetostratigraphic events; their calibration with astronomical cyclicity allowed a detailed chronostratigraphy to be established (Krijgsman et al., 1999a,b; Vai, 1997).

ii) the *Gessoso-solfifera Fm* (GS, Messinian), is made up of both primary and clastic, resedimented

evaporites with interbedded organic-rich shales, deposited during the evaporitic and post-evaporitic stages of the MSC;

iii) the *Colombacci Fm* (CO, Upper Messinian), consisting of mainly siliciclastic sediments derived from Apenninic sources, was deposited in both shallow and deep brackish or freshwater basins developed during the Lago Mare phase of the MSC. Based on a sharp vertical facies change, this unit has been subdivided by Roveri et al., (1998) in two informal units, the *Tetto fm.* (below) and the

Cusercoli fm. (above). The Cusercoli fm is characterized in marginal basins by coarser-grained lithology and by the occurrence of thin lacustrine micritic limestone beds (*colombacci* limestones).

iv) the **Argille Azzurre Fm** (AA, Lower Pliocene), is made up of relatively deep marine mudstones deposited in a series of more or less connected wedge top basins locally encasing fan-delta conglomerates and shelf to perched-basin turbiditic sandstone bodies and small, isolated carbonate platforms deposits (*Spungone*).

From a physical-stratigraphic point of view, the Tortonian to early Pliocene succession has been subdivided into three large-scale synthem separated by large-scale unconformities recording important phases of tectonic deformation of the Apennine orogenic wedge. From base to top they are the T_2 (Tortonian-Messinian) synthem, the *MP* (Messinian-Pliocene), and *LP* (Lower Pliocene) syntems (Roveri et al., 1998; Roveri et al., 2001; Roveri et al., 2003). These large-scale units only slightly differ from those defined by Ricci Lucchi (1986) and can be further subdivided into lower rank U.B.S.U. delimited by minor unconformities and flooding surfaces (Roveri et al., 1998), still having a regional significance. The more reliable key events for long-distance correlations are the Tortonian/Messinian boundary (7.2 My), the Miocene/Pliocene boundary (5.33 My) and a widespread ash-layer within the post-evaporitic Lago Mare deposits, which holds a radiometric age of 5.5. My (Odin et al., 1997). Messinian deposits fall in the T_2 (pre- and evaporitic stage) and *MP* (pos-evaporitic stage) syntems. Hereafter the successions developing in the western and eastern sectors will be described separately.

Western sector - The western area extend from the Sillaro valley to the West up to the Bidente valley to the East. These physiographic boundaries correspond to two important tectonic alignment, as previously reported. The Tortonian to Pliocene succession of this sector is particularly well exposed (Figs. 9, 10) along the Santerno valley which will be visited during the first day of the field trip. The second day will be devoted to important details from outstanding outcrops along the Sintria and Lamone valleys.

T_2 synthem (late Tortonian-Messinian) - The base of the T_2 synthem corresponds to an abrupt facies change within the Mamoso-arenacea Fm. In previous stages turbiditic deposits are characterized by tabular geometries and high lateral continuity, suggesting deposition by unconfined currents flowing parallel to basin axis (i.e. in a NW-SE direction) in a nearly flat sea-bottom.

In the T_2 synthem, sandstone bodies are coarser-grained and show a lower lateral continuity; the abundance of basal scours, cut and fill and water-




Fig. 9 – Stratigraphy of the western sector.




Fig. 10 – Panoramic view of the stratigraphic succession cropping out in the Santerno valley.

escape structures suggest deposition from flows strongly affected by topographic constriction (Mutti et al., 1999). This facies change has been usually related to a regional phase of tectonic uplift and basin narrowing (Ricci Lucchi, 1981, 1986).

In the western sector, this is well represented by the Fontanelice "channels", two turbiditic sandstone bodies confined within large-scale erosional depressions and separated by a mudstone horizon (Ricci Lucchi, 1968, 1969, 1975, 1981).

The larger and thicker upper channelized body is encased in upper Tortonian mudstones (*ghioli di letto*). In a section almost normal to paleocurrents, the basal erosional surface deepens toward the SW and a spectacular NE-ward onlap of sandstone fill can be observed. The sedimentary fill is composite, with basal conglomerate bars (pebble composition indicating an Alpine source, Ricci Lucchi, 1969),

and at least two distinct sandstone bodies in the upper part. The upper Fontanelice system is overlain by a mudstone unit with interbedded smaller-size turbiditic sandstone bodies, forming an overall thinning-upward sequence. In the Santerno river section the mudstone unit contains a slump body, up to 50 m thick, characterized by sandstone olistoliths, derived by minor turbiditic bodies. In turn, the mudstone unit is capped by a 40 m thick horizon of organic-rich *euxinic shales*. The euxinic shales, characterized by a well-developed lithologic cyclicity have been studied in detail for bio-magneto- and cyclostratigraphic reconstructions in several sections (Monte del Casino, Monte Tondo, Monticino, Fig. 11; Vigliotti, 1988; Negri and Vigliotti, 1997; Krijgsman et al., 1999a; Vai, 1997). The topmost cycles are characterized by the development of thin carbonate layers.




Fig. 11 – Panoramic view of the late Tortonian-Messinian succession in the Santerno valley (stop 1.4). Note the chaotic horizon below the euxinic shales and the deformed gypsum unit on top.



Fig. 12 – Facies 3 selenitic gypsum in a block of the Garisenda tower of Bologna.

The T₂ synthem is topped by the shallow-water evaporites of the Gessoso-solfifera Formation, forming a continuous belt, up to 150 m thick from the Sillaro to the Lamone valleys. Gypsum deposits have a strong cyclical organization, as first recognized by Vai and Ricci Lucchi, (1977) which counted up to 16 small-scale, decametric-thick, shallowing-upward cycles recording the progressive evaporation of shallow lagoons. These cycles have been later interpreted as controlled by periodic changes of orbital parameters (Vai, 1997; Krijgsman et al., 1999b).

The ideal complete cycle consists of the vertical superposition of 6 sedimentary facies (Fig. 12, 13, 14), five of them made up of gypsum and carbonates;




Fig. 13– The ideal depositional cycle of the Vena del Gesso evaporites (Vai and Ricci Lucchi, 1977)

it starts with basal organic-rich shales (facies 1), deposited in shallow lagoons below the wave base; these shales are overlain by gypsified limestone stromatolites (facies 2), massive autochthonous selenitic gypsum (facies 3) and well-stratified autochthonous and reworked selenitic gypsum (facies 4). The size of selenite crystals usually decreases upwards; this change is usually thought to be related to the progressive evaporative draw down and increasing salt concentration of the lagoon. The

upper part of the modal cycles is characterized by nodular gypsum, acicular gypsum, gypsarenites (facies 5) and chaotic selenite breccia (facies 6). These deposits are interpreted as related to subaerial exposure and erosion at the end of an evaporative draw-down cycle, with strong gypsum reworking by waves or torrential streams action (Vai and Ricci Lucchi, 1977).

Progressive upward decrease in thickness and different facies distribution within individual cycles




Fig. 14. Vena del Gesso evaporites. A) Santerno river bed. Slumped stromatolitic limestone (F2) and limestone breccia (“calcare di base”) embedded in the topmost pre-evaporite mudstones; b) Rio Sgarba Quarry, Borgo Tossignano. GS, cycle II: euoxic shales (F1), and calc-gypsum laminites enclosing meter-sized “palisade”-type crystals of stromatolite-bearing gypsum (F2); c) Montebello Quarry, Marecchia Valley. GS, cycle III: upright and well-packed “swallow-tail” selenite crystals (F3). The evaporite growth is interrupted by an intra-cycle dissolution surface; d) Monte Tondo Quarry, cycle VI, reworked selenite (facies F4); e). “Swallow-tail” selenite crystal (F3): an opaque stromatolite-bearing core is surrounded by a transparent diagenetic overgrowth; f) Monte Tondo Quarry, cycle VII, reworked small selenite crystals with nodules and lenses of whitish selenite.

can be observed; upper cycles are thinner and autochthonous evaporite facies are much less well developed than the reworked clastic facies.

This suggests the superposition of an overall “regressive” trend on the smaller-scale, higher-frequency cyclicity. This trend provides the guideline for a precise cycle-by-cycle correlation throughout the Vena del Gesso basin, and further to the SE in the Marecchia valley area (epi-Ligurian evaporites). For correlation purposes, the complete succession of 16 evaporite cycles has been subdivided into 2 thin “basal” cycles (facies 1 and 2 only), 3 thick “major” cycles (facies 5 missing), a very thick 6th cycle (complete sequence) and 10 thin “minor” cycles (complete sequences). The described longer-term trend culminates with a great erosional unconformity cutting the gypsum unit at its top and marking the base of the overlying Messinian-Pliocene synthem (MP).

MP synthem (Messinian-Pliocene) - An angular unconformity is associated to the erosional surface (Vai, 1988) marking the T₂/MP boundary, clearly indicating its tectonic nature. This surface preserves spectacular evidences of subaerial exposure (karstic dykes filled by continental deposits rich in mammal fauna were found in the Monticino section, near Brisighella; Costa et al., 1986; De Giuli et al., 1988). The erosion associated to the MP unconformity deepens toward the southeastern end of this sector (Marzeno valley), where the evaporitic and pre-evaporitic Messinian deposits are completely missing.

Deposits of this synthem belonging to the Colombacci Fm and Argille Azzurre Fm are very thin and separated by the Miocene/Pliocene boundary. The uppermost Messinian Colombacci Fm consists mainly of gray to varicolored clays containing an hypohaline faunal assemblage (*Melanopsis spp.*, *Linnocardium spp.*). Small lenses of sandstone and pebbly sandstone and calcareous conglomerate locally occur. At places, a white micritic layer is also found; locally named *colombaccio* it gives the name to the formation. The Miocene/Pliocene boundary is marked by a characteristic dark, organic-rich horizon, whose origin is still poorly understood. Through the boundary the typical abrupt transition from non marine to relatively deep-marine waters is observed; such change is witnessed by the rich and diversified planktonic assemblages of the basal Pliocene.

Lower Pliocene deposits of this synthem consist of a thin unit (only 2 m thick in the Santerno section, according to Colalongo et al., 1982 and Cremonini et al., 1969) of deep-marine mudstones draping the whole area. Detailed biostratigraphic studies carried out in the Santerno and Monticino sections show that a large hiatus is associated to the regional

unconformity (LP) marking the top of the MP synthem. The LP unconformity occurs in the upper part of the Gilbert chron and is related to the advance of the Apennine compressive front. This event is recorded in the Sillaro-Santerno area by the sudden appearance within deep marine clays of coarse, chaotic deposits (pebbly mudstones and debris flow) derived from the Ligurian units (Cremonini and Ricci Lucchi, 1982). The low thickness of the early Pliocene deposits of the MP synthem in this area is essentially related to the locally strong erosion associated to the LP unconformity.

Deformation of primary evaporites - The gypsum unit of this sector is characterized by extensional and compressional deformations (Marabini and Vai, 1985), with rotated blocks and shallow thrust faults, partially affecting also the top of the underlying *euxinic shales*. Most of these deformations emanate from a detachment surface in the upper part of the *euxinic shales* (Marabini and Vai, 1985) and are sealed by MP deposits of the Colombacci Fm.

From W to E the deformation shows different characteristics. To the west (Santerno-Sillaro sector), the gypsum unit, which is thinner and more discontinuous, shows both compressive and extensional deformations. Rotational listric faults affect the gypsum unit on the left bank of the Santerno river (see stop 1.2 Fig. 27), while further to the west (M. Penzola) shallow thrust faults are responsible for the vertical repetition of the lower gypsum cycles (stop 1.8). Traces of anhydritization (due to higher lithostatic loading during burial) and subsequent rehydration have been observed from M. Penzola to the westernmost edge (Sillaro-Santerno valleys, Roveri et al., 2003).

To the east (Sintria valley) the deformation is more severe. Here the gypsum unit forms a complex imbricate structure made of three or more SW verging thrust sheets (Marabini and Vai, 1985; see stop 2.2 Fig. 36).

Eastern sector - No obvious unconformities occur in this sector throughout the considered stratigraphic interval. The uppermost Marnoso-arenacea Fm. here consists (Fig. 15) of a thick pile of tabular turbiditic sandstone bodies, made up of thick-bedded, coarse to very coarse and pebbly sandstones, with frequent amalgamated beds and basal scours. Such deposits form the Savio turbidite system, a composite unit consisting of up to five sandstone bodies vertically arranged in an overall fining-upward sequence with a maximum thickness of some 200 m, cropping out discontinuously along the Savio valley. A direct genetic link with the channelized systems of the western sector (Fontanelice systems) has been recently suggested (Roveri et al., 2002; Roveri et al., 2003) on the base of facies and regional geologic

considerations.

As in the western sector, these sandstone bodies are overlain by a mudstone unit, here more developed and containing two chaotic bodies separated by an undisturbed muddy horizon, 400 m thick. The chaotic bodies are made up of both intra and extrabasinal deposits (Lucente et al., 2002); their thickness reaches some 300 m.

An organic-rich unit corresponding to the *euxinic shales* of the western area occurs above the chaotic horizon; here such unit is thicker (100 m) and show a less evident lithologic cyclicity, due to the higher terrigenous content. As in the western sector, the Tortonian/Messinian boundary lies at the base of this unit allowing reliable correlations.

The most striking difference with respect to the western sector is the total absence of primary, shallow-water evaporites. The Gessoso-solfifera Fm. consists here of a thick complex of resedimented evaporites made up of well-stratified, thin-bedded gypsum turbidites and huge olistostromes containing large blocks of primary selenitic gypsum. These evaporites were deposited in relatively deep-waters, well below wave-base, as suggested by the observed sedimentary structures (Parea and Ricci Lucchi, 1972; Ricci Lucchi, 1973; Roveri et al., 2001; Manzi, 2001). Interbedded mudstones have a moderate organic content, decreasing upward. Neither traces of subaerial exposure, nor an obvious cyclical pattern can be recognized.

Clastic gypsum deposits are overlain by a large thickness (up to 600 meters) of terrigenous non-marine deposits of the Colombacci Fm in turn followed by Pliocene deep-marine deposits. The best outcrops of upper Messinian and lower Pliocene deposits occur in two large synclines (Giaggiolo-Cella and Sapigno synclines in Figs. 7, 15), possibly corresponding to local Messinian depocenters developed above the orogenic wedge. The basal Pliocene deposits have here the same facies character of the western sector (epibathyal mudstones belonging to the Argille Azzurre Fm.) as well as the Miocene/Pliocene boundary, characterized by the typical black shale horizon.

Summarizing, the Messinian succession has an overall basinal character. In such a depositional context, a reliable stratigraphic framework can be based on the recognition of abrupt vertical facies changes, characterized by both compositional and/or grain-sizes changes in deep-water deposits. Such change would correspond to large-scale reorganization of basin geometry and drainage pattern related to main tectonic events.

Using this approach with the integration of bio- and magneto-stratigraphic data, the T₂ unconformity can be traced at the base of the Savio turbidite system. As for the MP unconformity, clearly developed in




Fig. 15 – Stratigraphy of the eastern sector.




Fig. 16 - Tortonian to late Messinian evolution of the Apennine foredeep basin (axial section) showing the longitudinal fragmentation due to the growth of the Forlì line - related structural high (Roveri et al., 2002).

the western sector, Roveri et al. (1998, 2001) and Manzi (2001) suggested that it could be traced into a correlative conformity at the base of the resedimented evaporitic complex of this area. According to this view, the Gessoso-solfifera Fm. would belong to the MP synthem, and a time equivalent of primary evaporites occurs within the local euxinic shales.

Preliminary bio-magneto-stratigraphic data (Manzi, 2001) pointed out the occurrence of a highly organic-rich, barren horizon, never observed in marginal, “Mediterranean-type” successions. This horizon would represent the deep-water counterpart of the primary evaporites.

The MP synthem can be subdivided into two units (p-ev₁ and p-ev₂, see Roveri et al., 2001; Fig. 4), separated by a minor unconformity marking important paleogeographic and structural changes.

The lower unit (p-ev₁) only occurs in structural depressions and bridges the final Messinian gap (Krijgsman, 1999a,b). The p-ev₁ unit consists of a basal complex of resedimented evaporites, overlain by a thick, monotonous succession of finely laminated mudstones and siltstones containing minor turbiditic sandstone bodies and showing an overall coarsening-and shallowing-upward trend related to the progressive and rapid basin infill (Tetto fm.); an ash layer, dated at ca. 5.5 My (Odin et al., 1997), represents an exceptional lithostratigraphic marker throughout the whole Apennine foredeep basin, allowing to recognize the p-ev₁ unit across different sub-basins (Bassetti et al., 1994). Sedimentary facies and vertical trends together with geometric

characteristics (the cross-section triangular shape and stratal convergence toward structural highs reconstructed through surface and subsurface data) indicate the syntectonic nature of this unit.

The upper unit (p-ev₂) is tabular and thickens in structural depressions but also overlies and seals all the previously uplifted and subaerially exposed areas. This unit consists of a cyclic alternation of coarse and fine-grained tabular lithosomes arranged in an overall fining-upward trend. The basic cyclicity records the periodical, climatically controlled, activation of fluvio-deltaic systems dominated by catastrophic floods (Roveri et al., 1998). 3 to 4 sedimentary cycles are normally observed in the upper unit; a similar, but less evident, cyclicity is observed in the upper part of the p-ev₁ unit. The overall fining-upward trend suggests a backstepping of fluvio-deltaic systems possibly related to the progressive basin enlargement.

Geological cross-sections clearly show the onlap of this unit against the Forlì structural high and the progressive upward decrease of thickness and grain-size of the fluvio-deltaic sediments. Paleocurrents and facies changes (Manzi, 1997; Roveri et al., 1998) show that the entry points of fluvio-deltaic systems feeding the eastern Romagna basins (extensively outcropping in the Giaggiolo-Cella syncline – day 4 of the field trip) were located along the Forlì line. Coarse-grained bodies are regularly interbedded with muddy units containing three limestone horizons (*colombacci*). As in the western sector the uppermost one lies just below the M/P boundary, here marked, like elsewhere in the Apennine foredeep by a characteristic dark, organic-rich horizon.

Messinian events and the tectonic evolution of the Apennine foredeep

The comparison and correlation of the western and eastern sectors of the Romagna Apennines (Fig. 7) allow to reconstruct the sedimentary evolution of the Apennine foredeep basin during the Messinian. The most important feature that clearly arise from field data of this as well as of the other sectors of the Apennine thrust belt, is the strong control of tectonically-derived topography on the areal distribution and vertical evolution of depositional systems. During the Messinian an important uplift phase affected the Apennine thrust belt, leading to the closure and fragmentation of the Marnoso-arenacea foredeep basin, to its accretion to the orogenic wedge and to the development of a new depocenter in a more external position, now buried under the Po Plain. This phase of tectonic deformation led to the emersion of the Apennine backbone and to the development of an embryonal Apenninic drainage system, as witnessed by the




Fig. 17 – Line drawing of a seismic profile showing the buried culmination of the Riolo anticline. See location in the geological map of Fig 24 (from Roveri et al., 2003).

abrupt change of sediment composition and dramatic increase of sedimentation rate. The most impressive fact is the close time and genetic relationship that can be recognized between the tectonic history of the area and the main Messinian events. The Mediterranean-type Messinian succession of the western sector developed above and was delimited by an uplifting anticline related to the Forlì line (Riolo anticline in Figs. 7 and 16). Seismic data allowed Roveri et al. (2003) to trace this structure in the subsurface and reconstruct a single ENE verging arcuate structure plunging to the west, of which the FL represents the eastern lateral ramp (Fig. 17). The continuous uplift of this structure starting from the late Tortonian (Fig. 18) was responsible for the observed sedimentary evolution of the area. The first evidence of the growth of the Riolo anticline was the creation of a small topographic relief elongated parallel to the basin axis leading to the progressive narrowing of the Marnoso-arenacea basin with consequent lateral confinement of turbiditic flows running from the NW to the SE. The larger volume flows, accelerated by this lateral constriction, increased their erosional power and cut large-scale erosional features (the Fontanelice “channels”); their sediment load was probably carried further down basin to feed the Savio turbidite system. The thick sandstone beds of “channel” fills can be considered as lobes formed in a confined basin by smaller volume flows. The anticline growth along the frontal and lateral ramp split the foredeep into two different subbasins: an uplifting basin to the west and a subsiding basin to the north and to the east. The former foredeep basin was gradually cut-off from coarse-grained turbiditic sediment input and a muddy slope developed in both sectors. This tectonically active slope was characterized by strong instability leading to huge sediment failures. The deeper and more subsiding character of the eastern sector is witnessed by the higher thickness of Tortonian to lower Messinian succession and by the higher

terigenous content of the local euxinic shales. Independent evidence for the uplift of the western sector during the late Tortonian and early Messinian comes from sediment rates and paleobathymetric reconstruction (Kouvenhoven et al., 1999; Van der Meulen et al., 1999); their combination suggests a sediment rate decrease and a concurrent shallowing upward trend.

The decrease of sedimentation rate is essentially due to the suppression of the terrigenous component related to the progressive cut-off of this uplifting area from turbidity currents. During the Messinian the role of the Riolo anticline is clearly defined. Primary, shallow-water evaporites deposited only in the western sector within a large, semiclosed wedge-top basin. The longer-term shallowing-upward trend superposed on the small-scale gypsum cycles suggests a gradual upward reduction in the amount of accommodation space created. The evaporites are cut by an erosion surface – the intra-Messinian unconformity – of inferred subaerial origin that can be traced to the SE on the outcropping culmination of the structural high associated to the Forlì line lateral ramp and to the NE above the buried anticline crest revealed by seismic and well data. The unconformity is associated to an angular discordance and formed during a paroxysmal phase of tectonic activity. The time elapsed by erosion in marginal uplifting areas is recorded in topographic depressions by a volume of sediment corresponding to the lower post-evaporitic unit occurring in the eastern sector. As a consequence, in the Apennine foredeep basin a large-scale tectonic pulse stopped primary evaporite deposition and caused the transition from hypersaline to hyposaline basin. Deep-water settings never experienced desiccation and evaporitic sediments were emplaced by gravity flows, forming a thick unit with tabular geometry and onlap terminations against basin margins.

After the emergence following the evaporitic phase, the FL structural high was sealed in the latest




Fig. 18 – Evolution of the Riolo anticline during the Upper Tortonian-Lower Pliocene time interval (from Roveri et al., 2003).

Messinian,; the Lago Mare deposits resting above the intra-Messinian unconformity in the western sector belong to the uppermost p-ev₂ unit, as indicated by the occurrence of the colombarci limestones. The basal Pliocene flooding occurred over an almost flat topography developed in a phase of slow and generalized subsidence and basin enlargement following the intra-Messinian tectonic pulse. This phase lasted the whole early Pliocene, during which fine-grained, deep-water sediments accumulated throughout the Apennine orogenic wedge. A new important tectonic event, occurred at 4.2 My (LP

unconformity), marking a significant propagation of the Apennine thrust front toward the Po plain area (Ricci Lucchi, 1986). During this phase a generalized NE-ward tilting of the western sector led to the present-day structural setting, with the northern culmination of the Riolo anticline buried by a 1500 m thick Plio-Pleistocene cover.

Hot Messinian topics

In this section, some of the “hottest” Messinian topics are briefly introduced. They derive from data and interpretations presented in this field trip and should stimulate and focus the discussion around the larger scale implications of the Apennine foredeep Messinian record.

1 - The onset of the MSC - The onset of the evaporitic stage is commonly associated to a sea-level fall whose amplitude is not well-constrained (100 m according to Clauzon et al., 2001). The transition from euxinic shales to selenitic gypsum observed in the Apennine foredeep marginal successions is apparently abrupt. However, no reliable paleodepth indicators occur in the upper part of the pre-evaporitic deposits, due to low oxygen concentrations at the sea-bottom. Moreover, the stromatolitic limestones marking the base of the evaporitic unit do not necessarily imply a shallow-water origin (i.e. the photic zone). On the other hand, the Lower Evaporites are affected by severe post-depositional deformations flattening out within the upper part of the euxinic shales that are characterized by abundant shear planes. The commonly envisaged purely tectonic origin of such deformations has been recently contested by Manzi, (2001;) and Roveri et al. (2003). Indeed, the reconstruction of the sedimentary and tectonic evolution of the western sector as essentially related to the uplift history of the Riolo anticline, also provide arguments for the Messinian development of a low-angle W-SW dipping paleoslope, corresponding to the southern, inner flank of the anticline. Renewed uplift during the intra-Messinian tectonic phase could have triggered large-scale gravity failures of the gypsum unit on a detachment surface within the euxinic shales, thus accounting for the observed SW verging thin-skinned deformation. According to the extent of translation along this paleoslope, gypsum is actually superposed to sediments deeper than those above its original substratum. A possible vertical displacement of 150-200 m has been calculated (Manzi, 2001). As a consequence, the “abrupt” vertical transition from euxinic shales to primary evaporites could be overestimated.

2 - Cyclicity and stacking patterns of Lower

Evaporites - Actually, many differences from the above described ideal cycle of primary evaporites can be observed, especially concerning the regular superposition of facies with decreasing size of selenite crystals (facies sequence 3-4-5) is concerned; in some cases, inverse facies trends or symmetric rhythms (4-3-4) occur and their significance has not been fully understood yet. Such anomalous facies sequences have been observed at the same stratigraphic level (i.e. in the same cycle) in other areas (Marecchia valley), suggesting external forcing factors on facies development.

The aggradational stacking pattern of the Lower Evaporites is a common feature across all the Mediterranean geodynamic settings, and necessarily implies a generalized subsidence or a sea-level rise. The latter hypothesis (Clauzon et al., 2001) would imply a larger water exchange with the Atlantic Ocean, as also supported by the evaporite isotopic composition (Flecker and Ellam, 1999). The vertical facies changes observed in the evaporitic cycles of the Apennine foredeep indicate that a longer-term shallowing-upward trend is superposed upon small scale cycles. This indicates a gradual upward reduction in the rate of space creation suggesting competition between sea-level rise and tectonic uplift; the latter was possibly heralding the strong intra-Messinian tectonic pulse.

3 - The tectonic vs. eustatic origin of the intra-Messinian unconformity - The transition between the Lower Evaporites and the Upper Evaporites or Lago Mare stages of the MSC is marked in all marginal basins of the Mediterranean area by the development of a large erosional surface (the intra-Messinian unconformity), associated to a hiatus of variable amplitude. The origin of this erosional surface is commonly related to an evaporative sea-level fall in excess of 1,000 meters implying desiccation of the deepest Mediterranean basins. Such catastrophic event led to the subaerial exposure of Mediterranean continental margins and to a huge fluvial rejuvenation with the incision of deep canyons in front of largest rivers (Nile, Rhone; Clauzon, 1973, 1982; Ryan, 1978; Ryan & Cita, 1978).

The tectonically enhanced nature of the intra-Messinian unconformity due to a supra-regional deformational phase has been commonly rejected. However, in many basins developed in different geodynamic settings (Apennine foredeep, Tyrrhenian basins, Tertiary Piedmont Basin, Sicily, Eastern Mediterranean, Western Mediterranean) this erosional surface is clearly associated to an angular unconformity.

This suggests an important phase of structural reorganization all along the Africa-Eurasia

collisional margin (Meulenkamp et al., 2000). Duggen et al., (2003) envisaged complex deep-crustal or mantle processes occurring between 6.3 and 4.8 My, to explain the abrupt changes in magma composition of the Alboran volcanic belt and the large uplift (1 km) of the African-Iberian margin required to close the marine connections between the Atlantic and the Mediterranean. In the Apennine foredeep, this unconformity is associated to the most important deformational phase since the early Miocene, as it marks the emersion of the Apennine chain. The consequences of such paleogeographic changes on climate have not been investigated yet.

4 - Resedimented evaporites: origin and significance

The term clastic evaporites was first used in the late 60's-early 70's to indicate some "gypsiferous sandstones" cropping out in the Laga basin (the southern depocenter of the Messinian Apennine foredeep). These deposits were often associated with the "balatino gypsum", a well-laminated alternation of gypsum and bituminous shales, usually interpreted as a deep-water primary evaporitic facies. The occurrence within the "balatino gypsum" of sedimentary structures interpretable as megaripples (Fanantello river, Schreiber, 1973; see stop 3.3; Fig. 19) was not adequately taken into consideration. Actually, due to the recognition of very indicative sedimentary structures (load casts, fluid-escape structures, bed gradation, cross-lamination and tractions-plus-fallout structures), almost all the "balatino" gypsum facies of the Apennines has clearly to be re-interpreted as deposited from turbidity currents. The "balatino" facies is part of a wide spectrum of mass-flow deposits ranging from olistostromes to low-density turbidity currents. As a matter of fact, this class of deposits should be dealt with like the siliciclastic gravity flow deposits. However, due to the strong diagenetic effects typically underwent by evaporitic rocks, their study is a very difficult task.

Most of the evaporites of the Apennine foredeep are actually clastic deposits derived from the dismantling of primary, *in situ* evaporites and resedimented through gravity processes into relatively deep waters, below the wave base (Parea and Ricci Lucchi, 1972; Ricci Lucchi, 1973; Manzi, 2001; Roveri et al., 2001; Manzi et al., submitted). Despite their common occurrence, they have been virtually ignored until recent times. Based on an integrated sedimentologic, petrographic and geochemical study, Manzi et al. (submitted), have proposed a new facies classification and a genetic model for resedimented evaporates, which implies a close similarity with siliciclastic turbidites. This sedimentologic interpretation is strictly linked with the geodynamic model, which provides some constraints for the




Fig. 19- Graded gypsarenite layers (Fanantello section, stop. 3.3). Each layer is composed by a light -dark couplet related to original grain-size. A) Note load structures at the base of thickest layer and long wavelength climbing ripples at the top of the coarser division. B) thick layer with two climbing ripple divisions showing opposite paleocurrents.

modalities of gypsum detritus production and the definition of trigger mechanisms for the initiation of flow.

One basic problem is the definition of original grain-size of clastic evaporites; in Manzi et al. (submitted) a siliciclastic-derived relationship between grain size and sedimentary structures is proposed in order to indirectly define this important parameter in diagenetically transformed deposits.

On the basis of this approach, several facies and facies associations have been recognized. Their areal distribution is in good agreement with the hypothesis of the removal of primary evaporites formed in shallow thrust-top basins through large submarine collapses and glides triggered by tectonic-induced gravitational instability and their down current transformations into high and low-density turbidity currents. This is well evident considering facies distribution along a NW-SE transect from the Forlì Line to the eastern Romagna and northern Marche basin. Of course, local tectonically induced topography provides further complication to the general picture.

The general poor to absent terrigenous component in resedimented evaporite unit could suggest a genesis from submarine collapses. Fluvial floods can be thus discarded, also considering the paleogeographic setting of the Messinian. Resedimented evaporites were accumulated at the very beginning of the intra-Messinian tectonic phase that led to the first emersion of the Apennine chain. A land area with a relatively well-developed fluvial drainage only formed at the end of this uplift phase, as witnessed by the composition of terrigenous sediments of the p-ev₁ and p-ev₂ units. This is also suggested by the vertical facies sequences observed in the resedimented evaporite unit; a clear bipartition very

often occurs, with a lower part made up of well-stratified, fine-grained deposits and by an upper part made up of disorganized, very coarse clastites, dominated by slumping and olistostromes often containing large blocks of lithified primary selenite.




Fig. 20 – Genetic and stratigraphic relationships between resedimented and primary evaporites (from Manzi, 2001).

This vertical organization could well reflect a sort of “progradation” of the system related to the ongoing tectonic deformation; however, we believe that it rather records the progressive uplift and consequent exhumation and denudation of primary evaporitic units formed above or around structural highs (Fig. 20). According to this view, basal fine-grained resedimented evaporites could derive from the erosion of the thinner-bedded, mainly reworked and unlithified uppermost cycles. The subsequent erosion of the early-cemented, thicker-bedded and coarser-grained primary evaporites lower cycles could produce olistostromes and debris flows. As a consequence, a sort of “inverted stratigraphy” is recorded by the resedimented evaporitic complex, which could follow-up with the muddy deposits of the p-ev1 unit deriving from the euxinic shales and underlying upper Tortonian slope mudstones. Resedimented evaporites could be a common feature in other Messinian basins of the Mediterranean area. Their correct recognition would represent a fundamental step for a better comprehension of the Messinian paleogeography and environmental changes. Of course, the recognition of other subaqueously deposited clastic evaporites associated to the intra-Messinian unconformity, would put into discussion the true extent of the Mediterranean deep desiccation.

5 - Gypsum diagenetic transformation and origin of sulphur mineralization - The Gessoso-solfifera Fm. of the Romagna and Marche Apennine contains sulfur deposits that were mined since Roman time until the '60s (Sciçli, 1972). The sulfur mineralization consists of discontinuous, thin lenses mostly restricted to the base of the evaporite formation and particularly at the contact of the sulfate layers with the basal organic-rich shales (euxinic shales). In some areas (Peticara) the mineralization is also present in higher stratigraphic levels into sulfate layers having particularly organic-rich shale interbeds. Sulfur-limestone layers appear to replace sulfate layers up to a few meters thick, but generally less than 1 m. The sulfur mean reach up to 40 %, but is generally around 12 %. The sulfur mineralization shows mostly nodular and breccia-veins textures and less common ribbon and massive textures. Native sulfur is associated to calcite, celestite, barite, bitumen and emissions of sulfuric acid and methane.

These characteristics indicate that elemental sulfur formed by sulfate reduction. Characteristics and timing of sulfur formation are unknown and detailed textural, petrographic and geochemical studies are at their initial stage. One of the significant aspects is that economically important sulfur concentrations are present only in the clastic sulfate evaporites of the

Romagna-Marche area that were transformed into anhydrite at burial diagenesis and not in the unaffected autochthonous selenite of the Vena del Gesso Basin. This suggests that early synsedimentary formation was negligible and that most sulfur originated by bacterial sulfate reduction triggered by hydrocarbons migrated from the associated organic-rich shales and marls. A thermochemical sulfate reduction origin for the sulfur can be excluded because the evaporite formation was buried at relatively shallow depth (~ 500-1000 m). Most sulfur deposition possibly occurred by oxidation of sulfuric acid by dissolved oxygen carried by groundwater during rehydration of anhydrite.

6 - The closure of the ‘Messinian gap’ and the onset of the Lago Mare event - In Messinian Mediterranean successions, the common stratigraphic gap between 5.50 and ~5.60 Ma is usually associated to the desiccation of deeper basins (Krijgsman et al., 1999b; Fig. 3). This gap does not allow a full understanding of the most important Messinian events: the switch from hypersaline to hyposaline waters leading to the Lago Mare stage. Causes, modality and timing of such dramatic paleogeographic change are obscure; the most commonly accepted explanation is the capture of Paratethyan waters by a lowered Mediterranean base-level.

As stated above, in the Apennine foredeep this gap only occurs above uplifted areas (e.g. the western sector of the Romagna Apennines); in deeper, subsiding basins the record is continuous and the hiatus being bridged by the lower part of the p-ev₁ unit, made up of resedimented evaporites and by a thick pile of terrigenous fine-grained sediments (Figs. 5, 21). This means that the Apennine successions have recorded such event and for this reason a detailed study of this interval is being carried out by a multidisciplinary research group. Preliminary observations and analysis carried out on a continuous core (Campea well, stop 4.1) led to the recognition of the typical Lago Mare ostracod, pollen and dinoflagellate assemblages immediately above the resedimented evaporite complex. This means that the typical Lago Mare hydrology was already established immediately after or during the deposition of the resedimented evaporite complex. Cores have also pointed out the occurrence of well-defined organic-rich horizons, which are being studied to assess their paleoenvironmental meaning and potential as regional lithostratigraphic markers.

7 - High-frequency Milankovitch-type climatic cyclicity in the Lago Mare deposits - Like pre-evaporitic deposits and lower Evaporites, the Lago Mare successions show a fairly well developed

cyclic organization that can be used to establish a high-resolution stratigraphy otherwise impossible to obtain in a non-marine unit encompassing a very short time span. According to Krijgsman et al. (1999a-b) the Lago Mare stage lasted more or less 270 Ky, 90 of them recorded by deep Mediterranean evaporites and by the basal p-ev₁ unit of the Apennine foredeep (Fig. 21). The bipartite character is a well recognizable but usually overlooked feature of the Lago Mare successions, not only in the Apennine foredeep but also in many other basins (see for example the Upper Evaporites and Arenazzolo Fm. of Sicily). Only the upper part of the lower unit, locally characterized by the deposition of shallow-water, sabkha-like evaporites (Sicily), shows a well-developed cyclical lithological pattern only in its upper part.

In the Apennine foredeep basins up to five cycles, defined on the base of the regular alternation of coarse and fine-grained lithosomes, can be recognized above the ash layer, which provide an independent, even if not particularly accurate, time calibration.

The upper p-ev₂ unit, which is widely recognized at a Mediterranean scale, is characterized by a well-developed high-frequency cyclical pattern superimposed upon an overall transgressive trend. These two characters represent good criteria for high-resolution long-range correlations; 3 to 4 main cycles have been traced from shallow to deep-water successions of the Apennine foredeep (Roveri et al., 1998; Ricci Lucchi et al., 2002), 4 cycles have been recognized in the uppermost Messinian deposits at Cyprus (Rouchy et al., 2001), 3-4 cycles in the latest




Fig. 21 – Correlation of Messinian Apennine foredeep succession with astronomically forced insolation cyclicity. Post-evaporitic deposits show a well-developed cyclic pattern especially in the upper unit (p-ev₂); p-ev₁ deposits, having a stronger syntectonic character, show a less-evident cyclicity, especially in the lower part.

Messinian Nile Delta deposits (Abu-Madi Fm., Dalla et al., 1997), 4 cycles in the fluvio-deltaic deposits of the Tertiary Piedmont Basin (Ghibaudo et al., 1985); 3 to 4 cycles are reported from a thick terrigenous Lago Mare succession unconformably overlying the Lower Evaporites in the Corvillo basin (Sicily; Keogh and Butler, 1999). As a consequence, a total of 8/9 cycles can be recognized in the 180 ky interval between the ash-layer and the Miocene/Pliocene boundary.

This is consistent with a precessionally-driven climatic origin of small-scale sedimentary cycles, like during the pre-evaporitic and evaporitic stages. By this way, the p-ev₂ unit would span a very short time interval of 60-80 Ky and a tentative correlation of its small-scale sedimentary cycles with the insolation curve is shown in Fig. 21. According to Roveri et al. (1998), the basic lithological cyclicality given by the vertical repetition of sharp-based coarse to fine-grained couplets, observed in both shallow, marginal basins developed above the orogenic wedge, as well as in the foredeep depocenters, would reflect the periodic activation of catastrophic flood-dominated fluvial systems carrying coarse sediments into the Lago Mare during dry intervals or at the transition between dry and wet periods (Fig. 22). During this phases (Fig. 5), small forestepping deltaic systems formed in shallow basins with low gradient shelves; in basins with reduced or steep shelves, fluvial floods could develop hyperpycnal flows able to carry their sediment load down basin forming small, turbidite systems of “low-efficiency” (*sensu* Mutti et al., 1999). Water carried with sediments to the basin contributed to rapidly rise the base-level and caused, together with a concomitant climate change toward wetter conditions, the deactivation of flood-dominated deltaic systems and their landward migration. During wet phases, base level reached its maximum but sediment input was drastically lowered for both the reduction of catchment’s area and the possibly lower erosion rate due to the increased vegetation cover. In this phase only fine-grained, thinly laminated mudstones accumulated into the basin. In some places, rhythmic alternations of light and dark mudstone horizons give a typical banded aspect to the deposit, probably representing a distal lacustrine facies. Thin-bedded, finely laminated limestone horizons (“colombacci”) are usually found in the upper half of the cycles associated to mudstone deposits. According to Bassetti et al. (submitted - see topic 8) they derive from inorganic precipitation of micrite-size crystals in a relatively deep-water environment, possibly anoxic and related to events of permanent water mass stratification events during periods of lake level maxima and associated peaks of reduced terrigenous input. However, the concomitant occurrence in the

same part of the cycles of dark, organic-rich clay horizons rich in specialized Mollusk assemblages (*Congerie*) and interpreted as palustrine deposits, suggests a far more complicated stratigraphic architecture and evolutionary trends, as tentatively illustrated in the model of Fig. 22. Palustrine horizons found at different levels within mudstone units are not compatible with a simple deepening upward cycle. According to their position within each cycle, the fine-grained lithosome could actually be subdivided into a lower transgressive and an upper regressive unit, the latter topped by a subaerial erosional surface produced during base level fall in dry phases. In other words, a regressive hemicycle could be preserved in the uppermost part of the cycles. However, neither prograding coastal deposits or clear evidence of paleosols have been clearly recognized so far.

An exception is represented by the uppermost cycle (Roveri et al., unpublished data), where the dark layer marking the Miocene/Pliocene boundary locally overlies an irregular surface with weak




Fig. 22 – Ideal evolution of uppermost Lago Mare cyclicity in the Apennine foredeep basins. Modified from Roveri et al. (1998).

pedogenetic traces (Botteghino section in the Giaggiolo-Cella syncline).

However, the possibly very short time spans encompassed by subaerial exposure phases within each cycle, could have been not sufficient for paleosol development; moreover, the erosion produced by catastrophic fluvial floods would have removed any trace. According to the model of Fig. 22, palustrine horizons could develop above the subaerial exposure surfaces during initial base-level rises; the *colombacci* limestones would represent a sort of condensed section marking the maximum retreat of clastic sediment sources. Detailed studies of these deposits is being carried out to better understand their organization and paleoenvironmental meaning; however, they bear very strong analogies with the Lago Mare cycles recently described at Cyprus (Rouchy et al., 2001) from the topmost Messinian succession.

8 - Origin of the colombacci limestones – As previously reported, the colombacci limestones are a typical lithofacies closely associated to basinal mudstones of the p-ev₂ unit. They are abiogenic and formed as inorganic precipitates from the surface waters of the Lago Mare. According to Bassetti et al. (submitted), their precipitation required the periodic supersaturation of the epilimnion, possibly driven by climatically-induced events of lake-water stratification. This carbonate lithofacies is quite common in Lago Mare deposits of the Mediterranean basins; geochemical composition provide good elements to derive the hydrological structure of the basins and their degree of connection. As a general rule isotopic compositions show negative ¹⁸O values, indicating freshwater conditions or at least a high degree of water dilution. The high-resolution stratigraphic model of the Apennine foredeep made it possible to compare isotopic compositions of colombacci limestones occurring in each cycle in different sub-basins and hence to verify possible areal gradients and stratigraphic trends. The results (Fig. 23; Bassetti et al., submitted) show that the two lower horizons have homogenous, strongly negative isotopic values; the uppermost horizon has less negative values, almost falling in the “normal marine” range. This could either indicate a strong concentration due to evaporation in restricted basins or, on the contrary, a true connection with the Ocean. The latter would be in good agreement with the aggradational stacking pattern of the p-ev₂ unit, suggesting a overall “transgressive” trend heralding the Zanclean flooding.

9 - Marine incursions during the Lago Mare stage? – Besides the above considerations based on isotopic composition of limestone horizons, occasional




Fig. 23 – Isotopic composition of colombacci limestones showing the clear evolution toward the quasi-marine values field, suggesting higher water concentration or true incipient connection with ocean waters (from Bassetti et al., submitted).

marine incursions in the Lago Mare deposits have been reported from different Mediterranean basins, especially on the base of nannoplankton assemblages (Blanc-Valleron et al., 1998; Pierre et al., 1998; Spezzaferri et al., 1998; Snel et al., 2001). Isotopic composition of the typical faunal associations (Keogh and Butler, 1999), suggest the possibility that during the final part of this stage a large water body was present in the Mediterranean. As far as these notes were written, no clear indications of marine incursions have been found in the Apennine foredeep Lago Mare successions. However, the commonly observed foraminifera associations made up small-size, dwarfed specimens need to be thoroughly investigated.

10 - The Mio/Pliocene boundary - The Mio/Pliocene boundary, marking the sudden return to fully marine conditions, is a synchronous event at a Mediterranean scale (Iaccarino et al., 1999). A peculiarly interesting feature is that the direct superposition of MP11 deposits above rocks older than the uppermost Messinian, has never been reported. MP11 sediments are always associated to Messinian Lago Mare deposits, as part of a longer-term transgressive sequence. This is particularly clear in the Apennine foredeep where such a transition occurs in a phase of tectonic quiescence and generalized subsidence that probably led to the almost complete submersion of the Apennine backbone, previously emerged with the intra-Messinian uplift. Significantly, in successions




Fig. 24 – The characteristic black layer underlying the sharp transition to open and deep marine conditions at the Miocene/Pliocene boundary. Riccò section (western Emilia Apennines; from Roveri and Manzi, in press).

deposited around topographic highs and characterized by the permanent activation of fan-delta systems, the Miocene/Pliocene transition appears to be very gradual and such boundary becomes very difficult to recognize (Roveri and Gennari, unpublished data). Elsewhere, a black, organic-rich layer always underlies the lowest deep-marine MP11 sediments (Fig. 24). Its nature and significance are not clear and its systematic study is currently being carried out. As previously reported (topic 7), in some cases this organic-rich horizon seems to be associated with a slightly erosional surface and weak paleosol traces.

According to the above mentioned possible occurrence at the end of the Messinian of a large Mediterranean non-marine water body and a higher degree of connection among the different sub-basins, the Miocene/Pliocene transition would be rather explained by a sudden hydrologic change associated to a bathymetric change of lower amplitude than usually thought. This would point to a less catastrophic character of the Zanclean flooding.

Field trip itinerary

Introduction to Day 1 and 2

These two days will be spent in an area between the Sillaro and the Lamone valleys (Fig 25). The Santerno Valley that will be visited during day 1 surely offers one of the most continuous and spectacular Neogene sections of the Northern Apennines and is particularly famous for the outcrops of the Miocene Marnoso-arenacea Fm, described in great detail in classical papers by Franco Ricci Lucchi and his co-workers in the late 60's and 70's. The local stratigraphic succession that can be observed spans in age from Early Serravallian to Pliocene. The Marnoso-arenacea Formation is overlain by the Messinian evaporites which, in turn,

are unconformably overlain by uppermost Messinian continental deposits and marine Pliocene-Pleistocene strata. The geology of the Santerno valley is relatively simple at the surface, where strata form a regular homoclinal gently dipping to the northeast, i.e. down the valley. Moving upstream from Imola, we drop down into progressively older stratigraphic levels. A packet of vertical to slightly overturned strata interrupts this regular bedding attitude at Coniale; these beds represent the northern limb of the ramp anticline associated to the M. Castellaccio thrust, a tectonic feature elongate in a NW-SE direction that can be traced as far as the Savio valley. In the Santerno valley, the core of the Coniale anticline is where the oldest stratigraphic level of the Marnoso-arenacea Fm is exposed. Further upstream, between S. Pellegrino and Firenzuola, the sand-rich Firenzuola system turbidites crop out in the southern flank of the Coniale anticline.

Day 1

Stop 1.1 - The reception center of the Vena del Gesso Natural Park, Tossignano. Depending on time availability and weather conditions, this optional stop will be devoted to visit a small museum exhibiting materials about nature, geology, mining activities, history and country life in the peculiar environment of the Vena del Gesso.

Stop 1.2 – Rocca di Tossignano. The ruins of the medieval castle on the hilltop are a wonderful place for a panoramic view over the Santerno valley and the Apennine foothills monocline (see cover photo and Fig. 26). From SW to NE, the geologist look can explore up-section the following stratigraphic units, whose details will be dealt with in the next stops:

- Marnoso-arenacea Fm, Fontanelice member: coarse sandstone and conglomerate bodies, composed of multiple or individual “channel” fills.




Fig. 25 – Geological map of the western sector




Fig. 26 – Stop 1.3. The spectacular Lower Evaporites outcrop (Rio Sgarba sect.), looking eastward from Tossignano.

- Marnoso-arenacea Fm., “ghiola di letto” unit: fine-grained thin-bedded turbidites and marls in an overall thinning- and fining-upward trend.
- Marnoso-arenacea Fm, “euxinic shales” unit: marls embedding cyclic intercalations of bituminous clays. The Tortonian-Messinian boundary is situated about 60 m below the unit top.
- Gessoso-solfifera Fm: it is composed of up to 16 evaporite cycles, subdivided into 2 “basal” ones (not visible), 3 “major” ones (grey selenite), a very thick

- 6th one (grey selenite and white clastic gypsum), and up to 10 “minor” ones (mostly white clastic gypsum). All 16 cycles crop out only in the eastern side of the Santerno valley (Rio Sgarba quarry, stop 1.4). In the western valley side (Fig. 27) a maximum of 15 cycles is found in the old Paradisa quarry, while 7 cycles crop out at M. Penzola, and only 4 ones crop out at Orzara. Near the Santerno river bed, this W-ward reduction of the evaporite succession is clearly explained by a late Messinian angular unconformity (Fig. 27), while further to the W a non-depositional hiatus is also possible. This alternative explanation would be in agreement with the overall W-ward thinning of the individual evaporite cycles from the Senio valley to the Sillaro valley (Fig. 26).
- Colombacci fm: a hard-to-see horizon of continental upper Messinian deposits (p-ev₂ sequence), which unconformably cover the Gessoso-solfifera Fm and the pre-evaporite shales. Thickness and facies of this fm are controlled by structural highs and lows of the evaporite unit (Fig. 28). For instance, conglomerates and sandstone pockets (facies H7-H8 in Fig. 29) are better represented within structural depressions.
- Argille Azzurre Fm: marine silty clays of early Pliocene age. Close to the western divide of the Santerno Valley, clays embed olistostromes and thick bodies of resedimented conglomerates, fed by epi-Ligurian fan-deltas. The well-stratified setting of all Pliocene units is clearly reflected in the landscape morphology (cover photo; Fig. 28).




Fig. 27. Stratigraphic columns of the Vena del Gesso in the western side of the Santerno valley (after Landuzzi, 1985): A= euxinic shales and stromatolitic limestones; B= euxinic shales, calc-gypsum laminites and gypsified stromatolites (facies 1-2); C= massive autochthonous selenite (facies 3); D= interbedded autochthonous and reworked selenite (facies 4); E= entirely reworked selenite (facies 4?); F= clastic, acicular and nodular gypsum (facies 5); G= chaotic gypsum in a clayey matrix (facies 6); H= post-evaporite deposits (see details in Fig.28).

Stop 1.3 - Tossignano, Resistance (II World War) Memorial Park. This panoramic view covers the GS from the Riva San Biagio cliff (Fig. 26) to the Rio Sgarba quarry. Basal, major and minor cycles can be easily distinguished and correlated. Two slump horizons embedded in the 8th and 10th evaporite cycles can be used as additional marker beds. The provenance of those slumps is still unknown, but might be an important constraint to the evolution of the Riolo Terme high (Fig. 25). Evaporites are offset by late Messinian normal faults belonging to a Graben structure symmetrical to the bedding planes




Fig. 28. Air photo of the Vena del Gesso in the western side of the Santerno Valley (M. Astorri). Normal faults and very small reverse faults offset the evaporite cycles, and are sealed by post-evaporite Messinian deposits, which mark a spectacular angular unconformity. The rotational offset of faulted blocks suggests gravity-driven normal faulting, gliding and thrusting on a shallow detachment level within the euxinic shales. Transport was to the S-SW.

(Fig. 30). Perhaps, those normal faults can be associated to gravity-failure structures, like those documented on the other side of the Santerno valley. **Topics:2.**

Stop 1.4 - Road Tossignano - Borgo Tossignano – Codrignano; deviations to the abandoned quarry in the Rio Sgarba valley, and to the old quarry at Paradisa, on the left bank of the Santerno river. Both sites were deeply investigated when the model of Vai and Ricci Lucchi (1973) was elaborated. The peculiar facies architecture of the modal cycle can be assessed in detail starting from the 6th cycle, which is the first and thickest example of a complete facies sequence. Smaller scale discrepancies with the basic model represent an interesting subject for debate. For instance, the complex and anomalous crystal size variation in the 3rd, 4th and 5th cycle (Fig. 27), where reverse gradation, dissolution surfaces (Fig. 14 c) and reworked selenite intervals are common. These intra-cycle features, possibly related to brine dilution episodes, have the same impressive correlation potential as the entire modal cycles. Indeed, they can be traced throughout the para-autochthonous Vena del Gesso basin, and from it to the semi-allochthonous evaporite basin of the Marecchia-Conca area (Montebello, Gesso, Sassofeltrio). Another basin-scale problem is represented by the sharp and strongly irregular erosion surface marking the onset of facies 5 in the 6th cycle. Some of the possible large-scale factors controlling evaporite deposition are summarized in the “hot Messinian topics” chapter.

Topics:2,7

Stop 1.5 - Road Borgo Tossignano - Fontanelice, Molino Campola. A panoramic view from the Santerno river bed to the western side of the valley allows framing a 50-60 m thick intraformational slump within the fine-grained closure facies of the MA (Fig. 31). The inferred slumping direction to the S-SW can be related to the late Tortonian nucleation




Fig. 29 Stratigraphic columns of the Colombacci fm (upper Messinian p-ev2 unit) in the western side of the Santerno Valley (after Landuzzi, 1985): C-G = GS facies 5-6 (see Fig. 27); H1 = limestone breccia with gypsum moulds; H2 = fine-grained clastic limestone; H3 = “varve”-like silty clays with fine-grained limestone lenses; H4 = laminated silty clays; H5 = variegated green and yellow clays; H6 = variegated green and red clays; H7 = poorly sorted calcareous conglomerates in small channel bodies; H8 = festoon-bedded sandstones in small channel bodies; H9 = fossil-rich mudstones; H10 = dark organic-rich pebbly clays; P = Lower Pliocene marine clays (Argille Azzurre Fm).




Fig. 30. Air photo of the Vena del Gesso in the eastern side of the Santerno Valley (M. Astorri). Longitudinal (FCN, FMA) and transverse (FCL) normal faults offset the evaporite cycles, and are sealed by post-evaporite Messinian deposits (p-ev2).

of the Riolo Terme high (Fig. 25).

Topics:2.

Stop 1.6 - Road Fontanelice – Gesso. Other panoramic views illustrate the closure facies and the earliest deformation features of the MA clastic wedge. The uppermost “channel” body of the Fontanelice member (Figs. 9, 10), the already described slump and the overall thinning- and fining-upward trend are coherent indications of foredeep fragmentation and basin narrowing.

Topics:1, 2

Stop 1.7 - Pieve di Gesso and Sassatello valley A panoramic view on the so-called “Sillaro line” portrays the syndepositional overthrust of the Liguride nappe in Messinian times (Fig. 32). Olistostromes coming from the Liguride and epi-Ligurian units have been continuously forerunning




Fig. 33. The Liguride nappe thrusting over the western Romagna para-autochthon: Messinian pre-evaporite (MA), evaporite (GS) and post-evaporite units are truncated atop by the overthrust surface, a gentle ramp dipping to the W. Olistostromes fed by Liguride and epi-Ligurian units occur both above and below the tectonic contact. Near Sassatello, a peculiar Messinian olistostrome is embedded in the p-ev2 unit of the para-autochthon.




Fig. 31. Closure facies of the Marnoso-arenacea Fm in the western side of the Santerno Valley: two sandstone olistoliths mark a slump horizon in a fining-upward succession of thin-bedded turbidites. Local slumping transport was possibly to the S-SW.

the nappe advancement. Many of them were tectonically incorporated in the nappe itself, while other were embedded in the autochthon succession. An example of the second type is the upper Messinian Sassatello olistostrome, enclosed in the para-autochthonous p-ev₂ unit. Its composition is characterized by lower-middle Miocene marls and typical Messinian carbonates, such as pre-evaporite and evaporite-derived limestones. A closer inspection of the olistostrome and the local Gessoso-solfifera Fm gives us the opportunity to discuss the diagenetic transformations of primary evaporites.

Topics:1, 2, 5.

Stop 1.8 - The "M. Penzola walk" We walk along the Santerno – Sellustra watershed from M. la Pieve to M. Penzola and Debolezza, then we go down to Casetta Gessi, and reach the nearest bridge on the Santerno river. (Fig. 28) The main subjects that are dealt with are the lower Pliocene clays and conglomerates, the angular unconformities in the post-evaporite p-ev₂ sequence and the late Messinian tectonics. The most attractive features of this itinerary are the M. Penzola thrust (Fig. 34), and a short but complete section of the Colombacci fm (Fig. 35, Fig. 29), featuring a very good outcrop of



Fig. 34. Monte Penzola, western side of the Santerno Valley: a thin-skinned thrust carries the 2nd evaporite cycle over the 5th one. The N-NE vergency of the thrust is contrary to the S-SW transport of all the Vena del Gesso gravity structures. This anomaly might result either from complex accommodation of gliding blocks, or from “real” tectonic thrusting.



Fig. 35. Northern slope of Monte Penzola: an about 5 m thick Colombari fm (see Fig. 29) separates the 6th cycle, facies 5 from the lower Pliocene clays. The typical dark layer marking the Miocene/Pliocene boundary is slightly offset by normal faults.

the Miocene/Pliocene boundary. Discussion is expected on the role of gravity vs. tectonics in the cortical deformation style of the evaporite unit.

Topics: 1, 2.

Day 2

Stop 2.1 - Visit to the reception centre of the Vena del Gesso Natural Park, Riolo Terme. This is another optional stop about the geologic, ecologic, historical and socio-economical importance of the Vena del Gesso.

Stop 2.2 - Road Riolo Terme - Casola Valsenio, stop at Borgo Rivola: the M. Tondo quarry and the adjoining Messinian outcrops. Monte Tondo and Monte del Casino are respectively situated on the eastern and western divide of the Senio valley. Actually, they are the two most important sampling sites for high-resolution cyclostratigraphy, magnetic calibration and astronomic dating of the main Messinian events in the Vena del Gesso basin.

Discussion will mainly focus on the cyclicity of the pre-evaporite unit. If possible, participants will be driven to the site of the M. Tondo section, which straddles the Tortonian-Messinian boundary and extends upwards to the GS base. The typical pre-evaporite Messinian is a dm-scale dark-light alternation of thin-laminated bituminous shales (anoxic levels), non-laminated organic-rich clays (disoxic levels) and common marly clays (oxic levels).

Topics: 1, 2.

Stop 2.3 – Zattaglia - Brisighella road, near La Torretta. In a panoramic view over the western side of the Sintria Valley (Fig. 36), the most prominent tectonic feature is a well-developed imbricate stack of SW-verging late Messinian thrust-sheets, which involve the GS and the upper part of the pre-evaporite succession. In the neighbourhood no similar structure is found, neither in the MA nor in the Argille Azzurre Fm. Indeed, back-thrusts are strictly confined to the evaporite unit. Another




Fig. 36 Monte Mauro and the western side of the Sintria Valley, from the S: the SW-verging thin-skinned thrusts that involve the evaporite unit can be explained by gravity-driven block-gliding, on a shallow detachment level within the euxinic shale unit (MA). As in the Santerno Valley example (Fig. 28), the initial effect of gliding was the development of SW-dipping normal faults dismembering the evaporite unit. During the subsequent gliding progression, the same faults were rotated and partly reactivated as thrusts (dashed lines).




Fig. 37. Monticino Quarry, Brisighella: the mining front represents the best possible exposure of the angular unconformity between the depositional sequences T2 and p-ev2. Some fractures sealed by the unconformity have been highlighted for their virtual relationship to the famous Vertebrate-bearing upper Messinian sedimentary dykes.

important character of this peculiar imbricate stack is that no one of the thrust-sheets has “roots” in the valley-floor, as no gypsum crops out in the river bed. This anomalous setting creates big problems for any cross-section reconstruction, if interpreted as a purely tectonic feature. Things get simpler if we consider a progressive gravity-driven slope failure. In the upper parts of the slope, the evaporite unit is extended and dismembered by rotational normal faults, while in its lower parts the faulted blocks keep gliding, and thrust over each other. This way, a stack of uprooted rock slices forms in the lower parts of the slope. Summarizing, by a merely structural point of view the back-thrusts cropping out in the Sintria and Lamone valleys are likely to have formed by gravitational failure and gliding. Other valid reasons are reviewed in the introduction of this Guide.

Topics: 1, 2

Stop 2.4 - Carne' Natural Park and/or Tanaccia cave entrance. A short off-road walk gives the participants the opportunity to see recent karstic morphologies superimposed on late Messinian thrust-faults and sub-vertical gypsum strata.

Topics: 1,2,3.

Stop 2.5 - Brisighella: the Monticino Sanctuary and the adjoining gypsum quarry. Deformation and emersion of the evaporite unit in late Messinian times are testified by an impressive angular unconformity between the Gessoso-solfifera Fm and the Colombacci Fm (Fig. 37). Subaerial erosion of gypsum is documented by karstic neptunian dikes sealed by post-evaporite Messinian deposits (p-ev₂). From 1985 to 1988, a very rich fauna of continental Vertebrates has been found in those paleo-karsts (Marabini & Vai, 1988). Panoramic and close views point out the present-day state of the quarry, which is being converted into an open-air geologic museum.

Topics: 1, 2, 3, 10


Stop 2.6 - Brisighella: participants are offered a leisure walk from the medieval castle to the Clock Tower, along faulted, vertical-bedded and karstified gypsum ridges.

Topics: 2.

Introduction to Day 3 and 4

In the Santerno section we examined a reduced, Mediterranean-type Messinian succession, developed above a structural high (Fig. 4). An abrupt facies and thickness change within the successions across the Forlì line that separates the marginal Vena del Gesso basin to the west from the eastern Romagna basins, is the best evidence of the strong structural control on Messinian deposition (Fig.7).

In the Savio valley area, we will examine Messinian deposits cropping out in the Sapigno and Giaggiolo-Cella synclines (see geological map of Fig. 38), two wedge-top basin bounded to the west by the Forlì thrust and filled up with more than 600 m of mainly post-evaporitic deposits. Shallow-water primary evaporites are lacking and are replaced by resedimented evaporites (turbiditic gypsarenites, olistostromes, olistoliths and breccias made up of selenitic and alabastrine gypsum). Sedimentary features (graded beds with erosional bases, traction plus fallout structures, sole marks, load casts, climbing ripples, coarse-grained beds deposited by hyperconcentrated flows, debris flow, small to large-scale intraformational slumpings) shows that these gypsum facies were deposited by gravity flows in a subaqueous setting. Siliciclastic deposits form the bulk of basin fill during the post-evaporitic stage. A very high sedimentation rate of more than 2mm/yr reflects the ongoing tectonic uplift and erosion of the Apennine chain following the intra-Messinian phase. Facies characteristics, paleoenvironmental meaning and cyclic stacking pattern of latest Messinian deposits will be also dealt with during these two days.






Fig. 39 – Stop 3.1. Suggested along basin correlation of upper Tortonian deposits between the Santerno and Savio valleys (datum plane: ~ the Tortonian/Messinian boundary; From Roveri et al., 2002).

Day 3

Stop 3.1 - Savio valley, panoramic views along the National road to Mercato Saraceno. The Upper Tortonian-Lower Messinian succession of the Savio valley: correlation with the Santerno valley (Fig. 39).
Topics: 1, 4.

Stop 3.2 – Road Sarsina - S.Agata Feltria, deviation to Maiano. Panoramic view of the resedimented evaporites complex of the Fanantello section (Fig. 40). Body geometry and stacking pattern.
Topics: 3, 4.

Stop 3.3 – Road Sarsina-S.Agata Feltria: the Rio Fanantello section (1h walk along the ‘Fanantello gorge’). The basal, stratified part of the resedimented evaporite complex crops out along the river bed offering a spectacular chance for a close view of facies characteristics of the *balatino* gypsum facies.
Topics: 4.

Stop 3.4 – Road Sapigno-Perticara. Panoramic view of the gypsum olistostrome and of the post-evaporitic Lago Mare succession of the Sapigno syncline.
Topics: 4, 6.

Stop 3.5 –Perticara: visit to the local Mine Museum.
Topic: 5.

Stop 3.6 - Road Perticara-Montepetra. Panoramic views and core data (Montepetra well) of the uppermost post-evaporitic (Lago Mare) deposits and the Miocene/Pliocene transition (Fig. 24).
Topics: 7, 8, 9, 10.

Stop 3.7 (optional) – Montepetra. Close inspection of the Montepetra “Lucina limestone”, a late Tortonian-early Messinian deep-water chemioherm related to cold fluid seepage at the sea-floor. Lucina limestones represent a classic topic for Apennine’s




Fig. 40– Stop 3.2. Panoramic view looking west from Maiano of the Messinian succession of the Sapigno syncline. Note the clear bipartite character of the resedimented evaporite complex overlying the local euxinic shales, also well detectable from slope morphology: the tabular gypsarenitic bodies of the lower part are overlain through an irregular surface by a huge chaotic gypsum complex.




Fig. 41. Facies and thickness distribution of the lowermost coarse-grained body of the Cusercoli fm (Corbara hzn)

geology; their time and space distribution suggest a strong relationships with the structural evolution of the Apennine thrust belt.

Day 4

Stop 4.1 - Road Piavola - Pian di Spino. Northern limb of the Giaggiolo-Cella syncline.

Panoramic and close views on general facies and physical-stratigraphic characteristics of the basal post-evaporitic succession; the transition to hyposaline conditions - the 5.5 Ma ash layer - sinsedimentary Messinian tectonics. Facies details of the p-ev₁ deposits in the cores of the Campea well.

Topics: 6.

Stop 4.2 – Road to Pieve di Rivoschio. Southern limb of the Giaggiolo-Cella syncline. Close view of an exceptionally thick gypsum layer (> 12 m) deposited by large-volume, high-density gravity flows. Like siliciclastic turbiditic beds, this gypsum layer has a lower coarser division with large-size mudstone clasts eroded and incorporated within the flow head, and an upper fine-grained laminated division with possible evidence of flow rebound. Gypsum recrystallization makes it difficult to assess the original textures, but general facies characteristics are still well recognizable.

Topics: 4.

Stop 4.3 – Road Pieve di Rivoschio – Voltre. Panoramic views of the post-evaporitic succession cropping out in the Giaggiolo-Cella syncline from its southern flank; general depositional characteristics, sedimentary trends and stratigraphic architecture of the p-ev₂ unit (Cusercoli fm). This unit is made up of three backstepping fluvio-deltaic coarse-grained bodies separated by lacustrine clays characterized by thin, laterally persistent limestone horizons (*colombacci limestones*).

Topics: 7

Stop 4.4 – Road Pieve di Rivoschio - Voltre (close view) - Corbara. Flood-dominated, fluvio-deltaic systems of the p-ev₂ unit. Close view of the coarse-grained deposits forming the base of small-scale cycles (Fig. 41). They consist of tabular bodies made up of amalgamated massive sandstones and pebbly sandstones, with scoured erosional bases and clay chips (depositional lobe), passing down current to hummocky cross-stratified sandstones and to thin bedded rippled sandstones; lobes are locally overlain by low-angle inclined conglomerates forming sigmoidal bars. Facies sequence has a regressive character and records the progradation in a shallow lacustrine basin of a small deltaic system dominated by catastrophic floods. Fine-grained deposits on top record the abandonment of the system due to a sudden decrease of flows volume.

Topics: 7

Stop 4.5 – Road Voltre - Pian di Spino. Anatomy of the upper Lago Mare cycles from outcrop and core data (Cà Blindana well). Sedimentology and palaeoenvironmental meaning of fine grained deposits of the p-ev₂ unit. Facies characteristics and origin of the “colombacci” limestones and of intervening black, organic-rich mudstones.

Topics: 7, 8, 9.

Stop 4.6 – Road Voltre - Cusercoli. Panoramic view showing the clear stacking pattern of the uppermost Messinian deposits. Climatic vs. tectonic control of high-frequency cyclicity; the latest Messinian ‘transgression’ and the Pliocene marine flooding.

Topics: 7

Stop 4.7 (optional) – Road Cusercoli-Meldola, Bidente valley. Close view of the Miocene/Pliocene boundary in the Buttafuoco section.

Topics: 10

Acknowledgments

The compilation of these notes wouldn't have been possible without the helpful work through the years of several students that we gratefully acknowledge: L. Capece, R. Gennari, A. Laffranchi, F. Rizzini, and M. Castellari.

The authors are also indebted to G. Papani, S. Iaccarino, R. Flecker, C. Schreiber, W. Schlager, E. Mutti, for their encouragement and helpful discussions about the different Messinian problems.

Reference cited

- Bassetti M.A., Ricci Lucchi F., Roveri M., 1994. Physical stratigraphy of the Messinian post-evaporitic deposits in Central-southern Marche area (Apennines, Central Italy). *Mem. Soc. Geol. It.* 48, 275-288.
- Bassetti M.A., 2000. Stratigraphy, sedimentology and paleogeography of upper Messinian ("Post-evaporitic") deposits in Marche area (Apennines, central Italy). *Mem. Sci. Geol.* 52/2, 319-349
- Blanc-Valleron M.M., Rouchy J.M., Pierre C., Badaut-Trauth D., Schuler, 1998. Evidence of Messinian nonmarine deposition at Site 968 (Cyprus lower slope), Proceedings of the Ocean Drilling Program: Scientific Results, Volume 160, Pages 437-446
- Boccaletti M., Calamita F., Deiana G., Gelati R., Massari F., Moratti G. and Ricci Lucchi F. 1990 Migrating foredeep-thrust belt system in northern Apennines and southern Alps. *Palaeogeography, Palaeoclimatology, Palaeoecology* 77, 3-14, Elsevier Science Publisher, Amsterdam.
- Capozzi R., Landuzzi A., Negri A., Vai G.B., 1991. Stili deformativi ed evoluzione Tettonica della successione neogenica romagnola. *Studi Geologici Camerti volume speciale* (1991/1), pp. 36-278.
- Castellarin A., Eva C.,iglia G, Vai G.B, with a contribute of Rabbi E., Pini G.A. and Crestana G. (1986) - Analisi strutturale del Fronte Appenninico Padano. *Giornale di Geologia*, 47 1/2, 47-75.
- Castellarin A., 2001. Alps-Appennines and Po Plain-frontal Apennines relations. In: Vai G.B. and Martini I.P., Eds, *Anatomy of an orogen: the Apennines and adjacent Mediterranean basins*. Kluwer Academic Publishers, Dordrecht-Boston-London, 177-196.
- Castellarin A., Pini G.A. with the contribute of Borsetti A.M. and Rabbi E., 1989. L'arco del Sillaro: la messa in posto delle Argille Scagliose al margine appenninico padano (Appennino bolognese). *Memorie della Società Geologica Italiana*, 39, 127-142.
- Clauzon G., 1973. The eustatic hypothesis and the pre-Pliocene cutting of the Rhone valley In Ryan W. B., Hsü K. J. Et al. Initial report of the Deep Sea Drilling Project. Washington XIII, 1251-1256.
- Clauzon G., 1982. Le canyon messinien du Rhone: une preuve decisive du "desiccated deep-basin model" (Hsü, Cita et Ryan, 1973): *Bulletin Societé Geologique de France* 24, 597-610.
- Clauzon G., Suc J.P., Gautier F., Berger A., Loutre M.F., 1996. Alternate interpretation of the Messinian salinity crisis: controversy resolved? *Geology* 24, 363-366.
- Clauzon G., Rubino J.L., Casero P., 2001. Regional modalities of the Messinian salinity crisis in the framework of a two phases model. 2nd EEDEN Workshop, Sabadell 2001, Abstract Book, 17-18.
- Colalongo M.L., Ricci Lucchi F., Guarnieri P, Mancini E., 1982. Il Plio-Pleistocene del Santerno (Appennino romagnolo), in Cremonini G., and Ricci Lucchi F., eds): *Guida alla Geologia del margine appenninico padano. Guide Geologiche Regionali della Società Geologica Italiana: Bologna, Società geologica italiana*, p. 161-166.
- Costa G.P., Colalongo M.L., De Giuli C., Marabini S., Masini F., Torre D. and Vai G.B, 1986. Latest Messinian Vertebrate fauna Preserved in a Paleokarst-neptunian dike setting (Brisighella, Northern Apennines): *Le grotte d'Italia*, v. 12, p. 221-235.
- Cremonini G., Elmi C., and Monesi A., 1969. Osservazioni geologiche e sedimentologiche su alcune sezioni plio-pleistoceniche dell'Appennino romagnolo: *Giornale di Geologia*, v. 35, p. 85-96.
- Cremonini G., and Ricci Lucchi F., 1982. *Guida alla Geologia del margine appenninico padano: Guide Geologiche Regionali, Società Geologica Italiana*, 248 pp.
- Dalla S., Harby H., Serazzi M., 1997. Hydrocarbon exploration in a complex incised valley fill: An example from the late Messinian Abu Madi Formation (Nile Delta Basin, Egypt). *The Leading Edge*, December 1997, 1819-1824.
- De Giuli C., Masini F. and Torre D., 1988, *The Mammal Fauna of the Monticino quarry*, in De Giuli, C., and Vai, G.B. eds., *Fossil Vertebrates in the Lamone valley, Romagna apennines*, International Workshop: Continental Faunas at the Mio-Pliocene Boundary, Faenza, March 28-31, 1988, *Guidebook*, p. 65-69.
- Duggen S., Hoernle K., Boogard P.v.d., Rupke L., Phipps

- Morgan J., 2003. Deep roots of the Messinian salinity crisis. *Nature*, 422, 602-606.
- Flecker R., Ellam, R.M., 1999. Distinguishing climatic and tectonic signals in the sedimentary succession of marginal basins using Sr isotops: an example of Messinian salinity crisis, Eastern Mediterranean. *Journal of the Geological Society*, London, 156, 847-854.
- Gelati R., Rogledi S., Rossi M., 1987. Significance of the Messinian unconformity-bounded sequences in the Apenninic margin of the Padan foreland basin, northern Italy (preliminary results). *Mem. Soc. Geol. It.* 39, 319-323.
- Ghibaudo G., Clari P., and Perello M., 1985. Litostratigrafia, sedimentologia ed evoluzione Tettonico-sedimentaria dei depositi miocenici del margine sud-orientale del Bacino Terziario Ligure-Piemontese (Valli Borbera, Scrivia e Lemme): *Bollettino della Società Geologica Italiana*, v. 104, p. 349-397.
- Hilgen F. J., Bissoli L., Iaccarino S., Krijgsman W., Meijer R., Negri A., Villa G., 2000. Integrated stratigraphy and astrochronology of the Messinian GSSP at Oued Akrech (Atlantic Morocco). *Earth and Planetary Science Letters*, 182, 237-251.
- Hsü K.J., Ryan W.B.F., Cita M.B., 1972. Late Miocene desiccation of the Mediterranean. *Nature*, 242, 240-244.
- Iaccarino S., M., Bossio A., 1999. Paleoenvironment of uppermost Messinian sequences in the western Mediterranean (sites 974, 975 and 978) In: Zahn R., Comas M.C. and Klaus A. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results 161*, 529-540.
- Iaccarino S., Papani G., 1980. Il Messiniano dell'Appennino Settentrionale dalla Val d'Arda alla Val Secchia: stratigrafia e rapporti con il substrato e il Pliocene. Vol. dedicato a S. Venzo, Univ. Studi di Parma, 15-46.
- Iaccarino S., Castradori D., Cita M. B., Di Stefano E., Gaboardi S., McKenzie J.A., Spezzaferri S., Sprovieri R., 1999. The Miocene/Pliocene boundary and the significance of the earliest Pliocene flooding in the Mediterranean. *Mem. Soc. Geol. It.* 54, 109-131.
- Keogh S. M. and Butler R.W.H., 1999. The Mediterranean water body in the late Messinian: interpreting the record from marginal basins on Sicily. *Journal of the Geological Society*, London, 156, 837-846.
- Kligfield R., 1979. The Northern Apennines as a collisional orogen: *American Journal of Science*, v. 279, p. 676-691.
- Kouwenhoven T.J., Seidenkrantz M.S. and Van der Zwaan G.J., 1999. Deep-water changes: the near-synchronous disappearance of a group of benthic foraminifera from the Late Miocene Mediterranean: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 152, p. 259-281.
- Krijgsman W., Hilgen F. J., Marabini S., Vai G. B., 1999a. New paleomagnetic and cyclostratigraphic age constraints on the Messinian of the Northern Apennines (Vena del Gesso Basin, Italy) *Mem. Soc. Geol. It.* 54, 25-33
- Krijgsman W., Hilgen F.J., Raffi I., Sierro F.J., Wilson D.S., 1999b. Chronology, causes and progression of the Messinian salinity crisis. *Nature* 400, 652-655.
- Lucente C.C., Manzi V., Ricci Lucchi F. and Roveri M., 2002. Did the Ligurian Sheet cover the whole thrust belt in Tuscany and Romagna Apennines? Some evidence from gravity emplaced deposits: *Memorie della Società Geologica Italiana*, 1, 393-398.
- Manzi V., 1997. Analisi di facies dei depositi messiniani post-evaporitici tra il Fiume Bidente ed il Fiume Savio: tra Corbara e Pieve di Rivoschio (Appennino romagnolo): unpublished Thesis, University of Bologna, 124 pp.
- Manzi V., 2001. Stratigrafia fisica, analisi sedimentologica microscopica e caratterizzazione magnetostratigrafica dei depositi connessi all'evento evaporitico del Messiniano (Formazione Gessoso-solfifera l.s.). Ph.D. thesis Earth Science Department, Bologna University, Italy.
- Marabini S., Vai G.B., 1985. Analisi di facies e macroTettonica della Vena del Gesso in Romagna. *Bollettino della Società Geologica Italiana*, v. 104, p. 21-42.
- Marabini S. and Vai. G.B., 2003. Marsili's and Aldrovandi's early studies on the gypsum geology of the Apennines. In Vai and Cavazza eds., *Fourth Centuries of the word Geology: Ulisse Aldrovandi 1603 in Bologna*. Minerva Edizioni.
- McKenzie J. A., 1999. From desert to deluge in the Mediterranean. *Nature* 400, 613-614.
- Meulenkamp J.E., Sissingh W. et al., 2000. Late Tortonian. In: S. Crasquin (Ed.), *Atlas Peri-Tethys, Palaeogeographical maps – Explanatory notes*. CCGM/CGMW, Paris, 195-201.
- Mutti, E., Tinterri, R., Remacha, E., Mavilla, N., Angella, S. and Fava, L., 1999. An introduction to the analysis of ancient turbidite basins from an outcrop perspective: AAPG Continuing Education Course Note Series, #39,
- Negri. A and Vigliotti L. 1997. Calcareous Nannofossil biostratigraphy and paleomagnetism of the Monte Tondo and Monte del Casino sections (Romagna Apennines, Italy) In (A. Montanari G.S. Odin and R. Coccioni, Eds): *Miocene Stratigraphy - An Integrated Approach*, Elsevier 478-491
- Odin G.S., Montanari A., and Coccioni R., 1997. Chronostratigraphy of Miocene stages: a proposal for the definition of precise boundaries, in (Montanari A., Odin G. S., and Coccioni R, eds): *Miocene Stratigraphy - An Integrated Approach*, Elsevier, p. 597-629.
- Parea G.C., Ricci Lucchi F., 1972. Resedimented evaporites in the periadriatic trough (upper Miocene, Italy) *Israel Journal of Earth-Sciences* 21, 125-141.
- Ricci Lucchi F., Bassetti M.A., Manzi V., Roveri M., 2002. Il Messiniano trent'anni dopo: eventi connessi alla crisi di salinità nell'avanfossa appenninica. *Studi. Geologici Camerti* 1, 127-142.
- Ricci Lucchi F., 1973. Resedimented evaporites: indicators of slope instability and deep-basins conditions in Peri-adriatic Messinian (Apennines foredeep, Italy). *Koninklijke Nederlandse Akademie Van Wetenschappen. Messinian Events in the Mediterranean. Geodynamics Scientific Report n° 7 on the colloquium held in Utrecht, March 2-4, 1973*, 142-149.
- Ricci Lucchi F., 1968. Chanellized deposits in the middle Miocene flysch of Romagna (Italy). *Giornale di Geologia*, 36, 203-282.
- Ricci Lucchi F., 1969. Recherches stratonomiques et sédimentologique sur le Flysch Miocène de la Romagne (Formation Marnoso-arenacea). *Proceedings of the 4th*

- session, Committee on Mediterranean Neogene Stratigraphy, *Giornale di Geologia*, 36, 163-198.
- Ricci Lucchi F., 1975. Depositional cycles in two turbidite formations of northern Apennines. *Journal of Sedimentary Petrology*, 45, 1-43.
- Ricci Lucchi F., 1981. The Miocene Marnoso-arenacea turbidites, Romagna and Umbria Apennines. In: Ricci Lucchi F., Ed., *Excursion Guidebook*, 2nd European Regional Meeting of International Association of Sedimentologists, 231-303.
- Ricci Lucchi F., 1986. The Oligocene to Recent foreland basins of the northern Apennines. *Spec. Publ. Int. Ass. Sediment.* 8, 105-139.
- Riding R., Braga J. C., Martin J. M., 2000. Late Miocene Mediterranean desiccation: topography significance of the "Salinity Crisis" erosion surface on-land in southeast Spain: Reply *Sedimentary Geology* 133, 175-184.
- Rossi M., Rogledi S., Barbacini G., Casadei D., Iaccarino S., Papani G., 2002. Tectono-stratigraphic architecture of the Messinian piggyback basin of the Northern Apennines: the Emilia folds in the Reggio-Modena area and comparison with the Lombardian and Romana sectors. *Boll. Soc. Geol. It.*, 1, 437-477.
- Rouchy J.M., Orszag-Sperber F., Blanc-Valleron M.M., Pierre C., Riviere M., Combourieu-Nebout N., Panayides I., 2001. Paleoenvironmental changes at the Messinian-Pliocene boundary in the eastern Mediterranean (southern Cyprus basins): significance of the Messinian Lago-Mare. *Sed. Geol.*, 145, 93-117.
- Roveri M., Bassetti M.A., Ricci Lucchi F., 2001. The Mediterranean Messinian salinity crisis: an Apennine foredeep perspective. *Sedimentary Geology* 140, 201-214.
- Roveri M., Manzi V., Bassetti M.A., Merini M., Ricci Lucchi F., 1998. Stratigraphy of the Messinian post-evaporitic stage in eastern-Romagna (northern Apennines, Italy). *Giornale di Geologia* 60, 119-142.
- Roveri M., Ricci Lucchi F., Lucente C. C., Manzi V., Mutti E., 2002. Stratigraphy, facies and basin fill history of the Marnoso-arenacea Formation. In: Mutti E., Ricci Lucchi F., Roveri M. Eds. *Revisiting turbidites of the Marnoso-arenacea Formation and their basin-margin equivalents: problems with classic models*. 64th EAGE Conference and Exhibition. *Excursion Guidebook*. Parma University and ENI – AGIP Division
- Roveri M., Manzi V., Ricci Lucchi F., Rogledi S., 2003. Sedimentary and tectonic evolution of the Vena del Gesso basin (Northern Apennines, Italy): Implications for the onset of the Messinian salinity crisis. *Geological Society of America Bulletin* 115, 387-405.
- Roveri M., Manzi V.. The Messinian salinity crisis: looking for a new paradigm? *Palaeogeography, Palaeoclimatology, Palaeoecology*. In press.
- Ruggieri G., 1967. The Miocene and later evolution of the Mediterranean Sea. Systematics association publication no. 7, *Aspects of the Tethyan biogeography*. Editors Adams C.G. and Ager D.V., 283-290, Palermo.
- Ryan W.B.F., Cita M. B., 1978. The nature and distribution of the Messinian erosional surface - indicators of a several-kilometers-deep Mediterranean in the Miocene. *Marine Geology* 27, 193-230.
- Ryan W.B.F., 1978. Messinian badlands on the southeastern margin of the Mediterranean Sea. *Marine Geology* 27, 349-363.
- Schreiber C.B., 1973. Survey of the physical features of the messinian chemical sediments Koninklijke Nederlandse Akademie Van Wetenschappen. *Messinian Events in the Mediterranean*. *Geodynamics Scientific Report n° 7* on the colloquium held in Utrecht, March 2-4, 1973. 101-110
- Scioli A., 1972. L'attività estrattiva e le risorse minerarie della regione Emilia Romagna. *Artioli Editore, Modena*, pp. 728.
- Snel E., Marunteanu M., Meulenkamp J.E., 2001. Late Miocene-Early Pliocene marine connections between the Atlantic/Mediterranean and the Paratethys. 2nd EEDEN Workshop, Sabadell 2001, *Abstract Book*, 69.
- Vai G.B., 1988. A field trip guide to the Romagna Apennine geology. The Lamone valley in De Giuli, C., and Vai, G.B. eds., *Fossil Vertebrates in the Lamone valley, Romagna apennines*, International Workshop: *Continental Faunas at the Mio-Pliocene Boundary*, Faenza, March 28-31, 1988, *Guidebook*, p. 7-37.
- Vai G.B., 1997. Cyclostratigraphic estimate of the Messinian stage duration: *in* Montanari, A., Odin, G.S., and Coccioni, R., eds.: *Miocene Stratigraphy - An Integrated Approach*: Amsterdam, Elsevier, p. 463-476.
- Vai G.B. and Ricci Lucchi F., 1977. Algal crusts, autochthonous and clastic gypsum in a cannibalistic evaporite basin: a case history from the Messinian of Northern Apennines.: *Sedimentology*, v. 24, p. 211-244.
- Vai G.B. and Martini I.P., 2001. *Anatomy of an orogen: the Apennines and adjacent Mediterranean*, Kluwer Academic Publishers, 632p.
- Van der Meulen M.J., Meulenkamp J.E., Wortel M.J.R., 1999. Lateral shifts of Apenninic foredeep depocentres reflecting detachment of subducted lithosphere. *Earth and Planetary Science Letters* 154, 230-219.
- Vigliotti L., 1988. Magnetostratigraphy of the Monticino section 1987 (Faenza, Italy), in De Giuli, C., and Vai, G.B. eds., *Fossil Vertebrates in the Lamone valley, Romagna apennines*, International Workshop: *Continental Faunas at the Mio-Pliocene Boundary*, Faenza, March 28-31, 1988, *Guidebook*, p. 61-62.
- Van Couvering J. A., Castratoro D., Cita M. B., Hilgen F. J., and Rio D., 2000. The base of the Zanclean Stage and of Pliocene series. *Episodes*, 23 (3), 179-187.

Back Cover:
field trip itinerary

FIELD TRIP MAP

